

OR4K5 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11998b**Specification**

OR4K5 Antibody (C-term) - Product Information

Application	WB, FC,E
Primary Accession	Q8NGD3
Other Accession	NP_001005483.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	36258
Antigen Region	284-313

OR4K5 Antibody (C-term) - Additional Information**Gene ID** 79317**Other Names**

Olfactory receptor 4K5, Olfactory receptor OR14-16, OR4K5

Target/Specificity

This OR4K5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 284-313 amino acids from the C-terminal region of human OR4K5.

Dilution

WB~~1:1000

FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR4K5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR4K5 Antibody (C-term) - Protein Information**Name** OR4K5**Function** Odorant receptor.


Cellular Location

Cell membrane; Multi-pass membrane protein.


OR4K5 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR4K5 Antibody (C-term) - Images

OR4K5 Antibody (C-term) (Cat. #AP11998b) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the OR4K5 antibody detected the OR4K5 protein (arrow).



OR4K5 Antibody (C-term) (Cat. #AP11998b) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

OR4K5 Antibody (C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR4K5 Antibody (C-term) - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)