PCDHGA8 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12016b

Specification

PCDHGA8 Antibody (C-term) - Product Information

Application	**IF, IHC-P, WB,E**
Primary Accession	Q9Y5G5
Other Accession	NP_114477.1
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**101480**
Antigen Region	**781-808**

PCDHGA8 Antibody (C-term) - Additional Information

Gene ID 9708

Other Names
Protocadherin gamma-A8, PCDH-gamma-A8, PCDHGA8, KIAA0327

Target/Specificity
This PCDHGA8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 781-808 amino acids from the C-terminal region of human PCDHGA8.

Dilution
IF~~1:10~50
IHC-P~~1:10~50
WB~~1:1000
E~~Use at an assay dependent concentration.

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
PCDHGA8 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PCDHGA8 Antibody (C-term) - Protein Information

Name PCDHGA8

Synonyms KIAA0327

Function Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.

Cellular Location
Cell membrane; Single-pass type I membrane protein

PCDHGA8 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

PCDHGA8 Antibody (C-term) - Images

Confocal immunofluorescent analysis of PCDHGA8 Antibody (C-term)(Cat#AP12016b) with U-251MG cell followed by Alexa Fluor 488-conjugated goat anti-rabbit lgG (green). DAPI was used to stain the cell nuclear (blue).

PCDHGA8 Antibody (C-term) (Cat. #AP12016b) western blot analysis in K562 cell line lysates (35ug/lane).This demonstrates the PCDHGA8 antibody detected the PCDHGA8 protein (arrow).

PCDHGA8 Antibody (C-term) (Cat. #AP12016b)immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of PCDHGA8 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

PCDHGA8 Antibody (C-term) - Background

This gene is a member of the protocadherin gamma gene cluster, one of three related clusters tandemly linked on chromosome five. These gene clusters have an immunoglobulin-like organization, suggesting that a novel mechanism may be involved in their regulation and expression. The gamma gene cluster includes 22 genes divided into 3 subfamilies. Subfamily A contains 12 genes, subfamily B contains 7 genes and 2 pseudogenes, and the more distantly related subfamily C contains 3 genes. The tandem array of 22 large, variable region exons are followed by a constant region, containing 3 exons shared by all genes in the cluster. Each variable region exon encodes the extracellular region, which includes 6 cadherin ectodomains and a transmembrane region. The constant region exons encode the common cytoplasmic region. These neural cadherin-like cell adhesion proteins most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been described for the gamma cluster genes.

PCDHGA8 Antibody (C-term) - References

Wu, Q., et al. Genome Res. 11(3):389-404(2001)
Nollet, F., et al. J. Mol. Biol. 299(3):551-572(2000)
Yagi, T., et al. Genes Dev. 14(10):1169-1180(2000)
Wu, Q., et al. Proc. Natl. Acad. Sci. U.S.A. 97(7):3124-3129(2000)
Wu, Q., et al. Cell 97(6):779-790(1999)