

PCDHB13 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12020c

Specification

PCDHB13 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	<u>Q9Y5F0</u>
Other Accession	<u>NP_061756.1</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	87552
Antigen Region	273-300

PCDHB13 Antibody (Center) - Additional Information

Gene ID 56123

Other Names Protocadherin beta-13, PCDH-beta-13, PCDHB13

Target/Specificity This PCDHB13 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 273-300 amino acids from the Central region of human PCDHB13.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions PCDHB13 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PCDHB13 Antibody (Center) - Protein Information

Name PCDHB13

Function Potential calcium-dependent cell-adhesion protein. May be involved in the



establishment and maintenance of specific neuronal connections in the brain.

Cellular Location Cell membrane; Single-pass type I membrane protein

PCDHB13 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

PCDHB13 Antibody (Center) - Images



PCDHB13 Antibody (Center) (Cat. #AP12020c) western blot analysis in MDA-MB435 cell line lysates (35ug/lane). This demonstrates the PCDHB13 antibody detected the PCDHB13 protein (arrow).

PCDHB13 Antibody (Center) - Background

This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gamma clusters, the transcripts from these genes are made up of only one large exon, not sharing common 3' exons as expected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment and function of specific



cell-cell neural connections.

PCDHB13 Antibody (Center) - References

Clark, H.F., et al. Genome Res. 13(10):2265-2270(2003) Frank, M., et al. Curr. Opin. Cell Biol. 14(5):557-562(2002) Vanhalst, K., et al. FEBS Lett. 495 (1-2), 120-125 (2001) : Wu, Q., et al. Genome Res. 11(3):389-404(2001) Nollet, F., et al. J. Mol. Biol. 299(3):551-572(2000)