

PAK6 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12164c

Specification

PAK6 Antibody (Center) - Product Information

Application	IF, IHC-P, WB,E
Primary Accession	O9NQJ5
Other Accession	NP_001122100.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	74869
Antigen Region	268-297

PAK6 Antibody (Center) - Additional Information

Gene ID 106821730;56924

Other Names

Serine/threonine-protein kinase PAK 6, PAK-5, p21-activated kinase 6, PAK-6, PAK6, PAK5

Target/Specificity

This PAK6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 268-297 amino acids from the Central region of human PAK6.

Dilution

IF~~1:10~50

IHC-P~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PAK6 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PAK6 Antibody (Center) - Protein Information

Name PAK6

Synonyms PAK5

Function Serine/threonine protein kinase that plays a role in the regulation of gene transcription. The kinase activity is induced by various effectors including AR or MAP2K6/MAPKK6. Phosphorylates the DNA-binding domain of androgen receptor/AR and thereby inhibits AR-mediated transcription. Also inhibits ESR1-mediated transcription. May play a role in cytoskeleton regulation by interacting with IQGAP1. May protect cells from apoptosis through phosphorylation of BAD.

Cellular Location

Cytoplasm. Nucleus. Note=Cotranslocates into nucleus with AR in response to androgen induction


Tissue Location

Selectively expressed in brain and testis, with lower levels in multiple tissues including prostate and breast


PAK6 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PAK6 Antibody (Center) - Images


Confocal immunofluorescent analysis of PAK6 Antibody (Center)(Cat#AP12164c) with U-251MG cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



PAK6 Antibody (Center) (Cat. #AP12164c) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the PAK6 antibody detected the PAK6 protein (arrow).



All lanes : Anti-PAK6 Antibody (Center) at 1:1000 dilution Lane 1: A431 whole cell lysate Lane 2: HT-29 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 75 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



PAK6 Antibody (Center) (Cat. #AP12164c) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of PAK6 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

PAK6 Antibody (Center) - Background

This gene encodes a member of the p21-activated kinase (PAK) family. The proteins of this family are Rac/Cdc42-associated Ste20-like Ser/Thr protein kinases, characterized by a highly conserved amino-terminal Cdc42/Rac interactive binding (CRIB) domain and a carboxyl-terminal kinase domain. PAK kinases are implicated in the regulation of a number of cellular processes, including cytoskeleton rearrangement, apoptosis and the MAP kinase signaling pathway. The protein encoded by this gene was found to interact with androgen receptor (AR), which is a steroid hormone-dependent transcription factor that is important for male sexual differentiation and development. This gene was found to be highly expressed in testis and prostate tissues and the encoded protein was shown to cotranslocate into the nucleus with AR in response to androgen. Alternatively spliced transcript variants encoding the same protein have been found for this gene. [provided by RefSeq].

PAK6 Antibody (Center) - References

Liu, C.Y., et al. Carcinogenesis 31(7):1259-1263(2010)
Guey, L.T., et al. Eur. Urol. 57(2):283-292(2010)
Lee, E.J., et al. Oncol. Res. 18(9):401-408(2010)
Hosgood, H.D. III, et al. Respir Med 103(12):1866-1870(2009)
Shen, M., et al. Environ. Mol. Mutagen. 50(4):285-290(2009)