

Mouse Parp9 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12195a

Specification

Mouse Parp9 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	O8CAS9
Other Accession	NP_084529
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	441-470

Mouse Parp9 Antibody (N-term) - Additional Information

Gene ID 80285

Other Names

Poly [ADP-ribose] polymerase 9, PARP-9, ADP-ribosyltransferase diphtheria toxin-like 9, ARTD9, B aggressive lymphoma protein homolog, Parp9, Bal

Target/Specificity

This Mouse Parp9 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 441-470 amino acids from the N-terminal region of mouse Parp9.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Parp9 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Parp9 Antibody (N-term) - Protein Information

Name Parp9

Synonyms Bal

Function ADP-ribosyltransferase which, in association with E3 ligase DTX3L, plays a role in DNA damage repair and in immune responses including interferon-mediated antiviral defenses (PubMed:[27796300](#)). Within the complex, enhances DTX3L E3 ligase activity which is further enhanced by PARP9 binding to poly(ADP-ribose) (By similarity). In addition, positively regulates DTXL3 protein levels (By similarity). In association with DTX3L and in presence of E1 and E2 enzymes, mediates NAD(+)-dependent mono-ADP-ribosylation of ubiquitin which prevents ubiquitin conjugation to substrates such as histones (By similarity). During DNA repair, PARP1 recruits PARP9/BAL1-DTX3L complex to DNA damage sites via PARP9 binding to ribosylated PARP1 (By similarity). Subsequent PARP1-dependent PARP9/BAL1-DTX3L-mediated ubiquitination promotes the rapid and specific recruitment of 53BP1/TP53BP1, UIMC1/RAP80, and BRCA1 to DNA damage sites (By similarity). In response to DNA damage, PARP9-DTX3L complex is required for efficient non-homologous end joining (NHEJ) but the complex function is restrained by PARP9 activity (By similarity). Dispensable for B-cell receptor (BCR) assembly through V(D)J recombination and class switch recombination (CSR) (PubMed:[28105679](#)). In macrophages, positively regulates pro-inflammatory cytokines production in response to IFNG stimulation by suppressing PARP14-mediated STAT1 ADP-ribosylation and thus promoting STAT1 phosphorylation (PubMed:[27796300](#)). Also suppresses PARP14-mediated STAT6 ADP-ribosylation (By similarity).

Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q8IXQ6}. Nucleus {ECO:0000250|UniProtKB:Q8IXQ6} Note=Shuttles between the nucleus and the cytosol. Translocates to the nucleus in response to IFNG or IFNB1 stimulation. Export to the cytosol depends on the interaction with DTX3L. Localizes at sites of DNA damage in a PARP1-dependent manner. {ECO:0000250|UniProtKB:Q8IXQ6}

Tissue Location

Highly expressed in the thymus and intestine (PubMed:18069692). Expressed in macrophages (PubMed:27796300)


Mouse Parp9 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mouse Parp9 Antibody (N-term) - Images





Parp9 Antibody (N-term) (Cat. #AP12195a) western blot analysis in mouse liver tissue lysates (35ug/lane). This demonstrates the Parp9 antibody detected the Parp9 protein (arrow).

Mouse Parp9 Antibody (N-term) - Background

PARP9 is a novel risk related gene that is expressed at higher levels in fatal high risk diffuse large B cell lymphomas.

Mouse Parp9 Antibody (N-term) - References

Hakme, A., et al. Dev. Dyn. 237(1):209-215(2008)