

**RPGR Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP12226b**

**Specification**

---

**RPGR Antibody (C-term) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">O92834</a>
Other Accession	<a href="#">NP_000319.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	744-772

**RPGR Antibody (C-term) - Additional Information**

**Gene ID** 6103

**Other Names**

X-linked retinitis pigmentosa GTPase regulator, RPGR, RP3, XLRP3

**Target/Specificity**

This RPGR antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 744-772 amino acids of human RPGR.

**Dilution**

WB~~1:1000

IHC-P~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

RPGR Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**RPGR Antibody (C-term) - Protein Information**

**Name** RPGR

**Synonyms** RP3, XLRP3

**Function** Could be a guanine-nucleotide releasing factor. Plays a role in ciliogenesis. Probably regulates cilia formation by regulating actin stress filaments and cell contractility. Plays an important role in photoreceptor integrity. May play a critical role in spermatogenesis and in intraflagellar transport processes (By similarity). May be involved in microtubule organization and regulation of transport in primary cilia.

#### Cellular Location

Cytoplasm, cytoskeleton, flagellum axoneme {ECO:0000250|UniProtKB:Q9R0X5}. Golgi apparatus. Cell projection, cilium {ECO:0000250|UniProtKB:Q9R0X5}. Note=In the retinal photoreceptor cell layer, localizes at the connecting cilium (By similarity). Colocalizes with WHRN in the photoreceptor connecting cilium (By similarity) Colocalizes with CEP290 in the photoreceptor connecting cilium (By similarity). Colocalizes with RPGRIP1 in the photoreceptor connecting cilium (By similarity). {ECO:0000250|UniProtKB:Q9R0X5}

#### Tissue Location


Heart, brain, placenta, lung, liver, muscle, kidney, retina, pancreas and fetal retinal pigment epithelium. Isoform 3 is found only in the retina. Colocalizes with RPGRIP1 in the outer segment of rod photoreceptors and cone outer segments

#### RPGR Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### RPGR Antibody (C-term) - Images



RPGR Antibody (C-term) (Cat. #AP12226b) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the RPGR antibody detected the RPGR protein (arrow).



Anti-RPGR Antibody (C-term) at 1:1000 dilution + human liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 113 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



RPGR Antibody (C-term) (Cat. #AP12226b) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of RPGR Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

### **RPGR Antibody (C-term) - Background**

This gene encodes a protein with a series of six RCC1-like domains (RLDs), characteristic of the highly conserved guanine nucleotide exchange factors. The encoded protein is found in the Golgi body and interacts with RPGRIP1. This protein localizes to the outer segment of rod photoreceptors and is essential for their viability. Mutations in this gene have been associated with X-linked retinitis pigmentosa (XLRP). Multiple alternatively spliced transcript variants that encode different isoforms of this gene have been reported, but the full-length nature of only some have been determined.

### **RPGR Antibody (C-term) - References**

Clark, G.R., et al. Ophthalmology 117(11):2169-2177(2010)  
Schmid, F., et al. Invest. Ophthalmol. Vis. Sci. 51(3):1628-1635(2010)  
Ji, Y., et al. Curr. Eye Res. 35(1):73-79(2010)  
Sheng, X., et al. Mol. Vis. 16, 1620-1628 (2010) :  
Murga-Zamalloa, C.A., et al. J. Genet. 88(4):399-407(2009)