

LIPC Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12283c

Specification

LIPC Antibody (Center) - Product Information

Application	FC, IHC-P, WB,E
Primary Accession	P11150
Other Accession	NP_000227.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	55914
Antigen Region	310-338

LIPC Antibody (Center) - Additional Information

Gene ID 3990

Other Names

Hepatic triacylglycerol lipase, HL, Hepatic lipase, Lipase member C, LIPC, HTGL

Target/Specificity

This LIPC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 310-338 amino acids from the Central region of human LIPC.

Dilution

FC~~1:10~50

IHC-P~~1:10~50

WB~~1:500

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

LIPC Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

LIPC Antibody (Center) - Protein Information

Name LIPC

Synonyms HTGL

Function Catalyzes the hydrolysis of triglycerides and phospholipids present in circulating plasma lipoproteins, including chylomicrons, intermediate density lipoproteins (IDL), low density lipoproteins (LDL) of large size and high density lipoproteins (HDL), releasing free fatty acids (FFA) and smaller lipoprotein particles (PubMed:[12032167](#), PubMed:[26193433](#), PubMed:[7592706](#), PubMed:[8798474](#)). Also exhibits lysophospholipase activity (By similarity). Can hydrolyze both neutral lipid and phospholipid substrates but shows a greater binding affinity for neutral lipid substrates than phospholipid substrates (By similarity). In native LDL, preferentially hydrolyzes the phosphatidylcholine species containing polyunsaturated fatty acids at sn-2 position (PubMed:[26193433](#)).

Cellular Location


Secreted.

LIPC Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


LIPC Antibody (Center) - Images



Anti-LIPC Antibody (Center) at 1:500 dilution + HT-29 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 56 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



LIPC Antibody (Center) (Cat. #AP12283c) immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of LIPC Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



LIPC Antibody (Center) (Cat. #AP12283c) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

LIPC Antibody (Center) - Background

LIPC encodes hepatic triglyceride lipase, which is expressed in liver. LIPC has the dual functions of triglyceride hydrolase and ligand/bridging factor for receptor-mediated lipoprotein uptake.

LIPC Antibody (Center) - References

Reynolds, R., et al. Ophthalmology 117(10):1989-1995(2010)
Jablonski, K.A., et al. Diabetes 59(10):2672-2681(2010)
Hu, M., et al. Pharmacogenet. Genomics 20(10):634-637(2010)
Romero, R., et al. Am. J. Obstet. Gynecol. 203 (4), 361 (2010) :
Kashani Farid, M.A., et al. Lipids Health Dis 9, 96 (2010) :