

EVPLL Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12312c**Specification**

EVPLL Antibody (Center) - Product Information

Application	IHC-P, WB,E
Primary Accession	A8MZ36
Other Accession	NP_001138599.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	33953
Antigen Region	118-146

EVPLL Antibody (Center) - Additional Information**Gene ID** 645027**Other Names**

Envoplakin-like protein, EVPLL

Target/Specificity

This EVPLL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 118-146 amino acids from the Central region of human EVPLL.

Dilution

IHC-P~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

EVPLL Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.


EVPLL Antibody (Center) - Protein Information**Name** EVPLL

EVPLL Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EVPLL Antibody (Center) - Images



Anti-EVPLL Antibody (Center) at 1:1000 dilution + MCF-7 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 34 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



EVPLL Antibody (Center) (Cat. #AP12312c) immunohistochemistry analysis in formalin fixed and

paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of EVPLL Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

EVPLL Antibody (Center) - References

Strausberg, R.L., et al. Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903(2002)