

DCC Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12334c

Specification

DCC Antibody (Center) - Product Information

Application	IHC-P, WB,E
Primary Accession	P43146
Other Accession	Q63155 , P70211 , NP_005206.2
Reactivity	Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	158457
Antigen Region	643-670

DCC Antibody (Center) - Additional Information

Gene ID 1630

Other Names

Netrin receptor DCC, Colorectal cancer suppressor, Immunoglobulin superfamily DCC subclass member 1, Tumor suppressor protein DCC, DCC, IGDCC1

Target/Specificity

This DCC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 643-670 amino acids from the Central region of human DCC.

Dilution

IHC-P~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DCC Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

DCC Antibody (Center) - Protein Information

Name DCC

Synonyms IGDCC1

Function Receptor for netrin required for axon guidance. Mediates axon attraction of neuronal growth cones in the developing nervous system upon ligand binding. Its association with UNC5 proteins may trigger signaling for axon repulsion. It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand. Implicated as a tumor suppressor gene.

Cellular Location

Membrane; Single-pass type I membrane protein.

Tissue Location


Found in axons of the central and peripheral nervous system and in differentiated cell types of the intestine. Not expressed in colorectal tumor cells that lost their capacity to differentiate into mucus producing cells.

DCC Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DCC Antibody (Center) - Images



DCC Antibody (Center) (Cat. #AP12334c) western blot analysis in mouse heart tissue lysates (35ug/lane). This demonstrates the DCC antibody detected the DCC protein (arrow).



DCC Antibody (Center) (Cat. #AP12334c) immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of DCC Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

DCC Antibody (Center) - Background

This gene encodes a netrin 1 receptor. The transmembrane protein is a member of the immunoglobulin superfamily of cell adhesion molecules, and mediates axon guidance of neuronal growth cones towards sources of netrin 1 ligand. The cytoplasmic tail interacts with the tyrosine kinases Src and focal adhesion kinase (FAK, also known as PTK2) to mediate axon attraction. The protein partially localizes to lipid rafts, and induces apoptosis in the absence of ligand. The protein functions as a tumor suppressor, and is frequently mutated or downregulated in colorectal cancer and esophageal carcinoma.

DCC Antibody (Center) - References

Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
Srour, M., et al. Science 328 (5978), 592 (2010) :
Docherty, S.J., et al. BMC Genet. 11, 61 (2010) :
Li, W., et al. Nat. Neurosci. 7(11):1213-1221(2004)