

NBPF5 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12364b

Specification

NBPF5 Antibody (C-term) - Product Information

Application WB, FC,E
Primary Accession Q86XG9

Other Accession <u>Q5VWK0</u>, <u>Q96M43</u>, <u>XP_001714524.1</u>

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Human
Rabbit
Polyclonal
Rabbit IgG
A0546
Action Region

263-291

NBPF5 Antibody (C-term) - Additional Information

Other Names

Putative neuroblastoma breakpoint family member 5, NBPF5P, NBPF5

Target/Specificity

This NBPF5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 263-291 amino acids from the C-terminal region of human NBPF5.

Dilution

WB~~1:1000 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NBPF5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NBPF5 Antibody (C-term) - Protein Information

Name NBPF5P

Synonyms NBPF5

Cellular Location



Cytoplasm.


Tissue LocationExpressed in brain and medulla.

NBPF5 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

NBPF5 Antibody (C-term) - Images



NBPF5 Antibody (C-term) (Cat. #AP12364b) western blot analysis in Jurkat cell line lysates (35ug/lane). This demonstrates the NBPF5 antibody detected the NBPF5 protein (arrow).







Tel: 858.875.1900 Fax: 858.875.1999

NBPF5 Antibody (C-term) (Cat. #AP12364b) flow cytometric analysis of Jurkat cells (right compared to a negative control cell (left histogram).FITC-conjugated donkey-anti-rabbit secondary antibodies were used for the analysis.