

DNAJB6 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12493c**Specification**

DNAJB6 Antibody (Center) - Product Information

Application	FC, IHC-P, WB,E
Primary Accession	O75190
Other Accession	NP_490647.1 , NP_005485.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	36087
Antigen Region	218-247

DNAJB6 Antibody (Center) - Additional Information**Gene ID** 10049**Other Names**

DnaJ homolog subfamily B member 6, HHDJ1, Heat shock protein J2, HSJ-2, MRJ, MSJ-1, DNAJB6, HSJ2, MRJ, MSJ1

Target/Specificity

This DNAJB6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 218-247 amino acids from the Central region of human DNAJB6.

Dilution

FC~~1:10~50

IHC-P~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DNAJB6 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

DNAJB6 Antibody (Center) - Protein Information

Name DNAJB6**Synonyms** HSJ2, MRJ, MSJ1

Function Has a stimulatory effect on the ATPase activity of HSP70 in a dose-dependent and time-dependent manner and hence acts as a co- chaperone of HSP70 (PubMed:[10954706](#), PubMed:[28233300](#)). Plays an indispensable role in the organization of KRT8/KRT18 filaments (PubMed:[10954706](#)). Acts as an endogenous molecular chaperone for neuronal proteins including huntingtin (PubMed:[11896048](#), PubMed:[22366786](#)). Suppresses aggregation and toxicity of polyglutamine- containing, aggregation-prone proteins (PubMed:[20159555](#), PubMed:[22366786](#)). Also reduces cellular toxicity and caspase-3 activity (PubMed:[11896048](#)).

Cellular Location

Cytoplasm, perinuclear region. Nucleus Cytoplasm, myofibril, sarcomere, Z line


Tissue Location

Widely expressed. Highest levels in testis and brain, and lower levels in heart, spleen, intestine, ovary, placenta, lung, kidney, pancreas, thymus, prostate, skeletal muscle, liver and leukocytes. In testis, expressed in germ cells in the earlier stages of differentiation pathway as well as in spermatids. In brain, expressed at a higher level in hippocampus and thalamus and a lower level in amygdala, substantia nigra, corpus callosum and caudate nucleus


DNAJB6 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DNAJB6 Antibody (Center) - Images

Western blot analysis of DNAJB6 Antibody (Center) Pab (Cat. #AP12493c) pre-incubated without (lane 1) and with (lane 2) blocking peptide in HeLa cell line lysate. DNAJB6 Antibody (Center) (arrow) was detected using the purified Pab.



DNAJB6 Antibody (Center) (Cat. #AP12493c) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of DNAJB6 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



DNAJB6 Antibody (Center) (Cat. #AP12493c) flow cytometric analysis of HeLa cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

DNAJB6 Antibody (Center) - Background

This gene encodes a member of the DNAJ protein family. DNAJ family members are characterized by a highly conserved amino acid stretch called the 'J-domain' and function as one of the two major classes of molecular chaperones involved in a wide range of cellular events, such as protein folding and oligomeric protein complex assembly. This family member may also play a role in polyglutamine aggregation in specific neurons. Alternative splicing of this gene results in multiple transcript variants; however, not all variants have been fully described.

DNAJB6 Antibody (Center) - References

Mitra, A., et al. J. Biol. Chem. 285(32):24686-24694(2010)
Edo De Bock, C., et al. Int. J. Oncol. 36(5):1155-1163(2010)
Hageman, J., et al. Mol. Cell 37(3):355-369(2010)
Lowe, J.K., et al. PLoS Genet. 5 (2), E1000365 (2009) :
Dey, S., et al. Mol. Cell. Biochem. 322 (1-2), 137-142 (2009) :