

NR2C2AP Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12673A**Specification**

NR2C2AP Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q86WQ0
Other Accession	Q3TV70 , Q58DU8 , NP_795361.1
Reactivity	Human
Predicted	Bovine, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	15876
Antigen Region	8-37

NR2C2AP Antibody (N-term) - Additional Information**Gene ID** 126382**Other Names**

Nuclear receptor 2C2-associated protein, TR4 orphan receptor-associated 16 kDa protein, NR2C2AP, TRA16

Target/Specificity

This NR2C2AP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 8-37 amino acids from the N-terminal region of human NR2C2AP.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NR2C2AP Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NR2C2AP Antibody (N-term) - Protein Information**Name** NR2C2AP

Synonyms TRA16

Function May act as a repressor of NR2C2-mediated transactivation by suppressing the binding between NR2C2/TR4 and the TR4-response element in target genes.

Cellular Location

Nucleus.


Tissue Location

Expressed in all tissues examined, with highest expression in heart, skeletal muscle and pancreas

NR2C2AP Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NR2C2AP Antibody (N-term) - Images

NR2C2AP Antibody (N-term) (Cat. #AP12673a) western blot analysis in Y79 cell line lysates (35ug/lane). This demonstrates the NR2C2AP antibody detected the NR2C2AP protein (arrow).

NR2C2AP Antibody (N-term) - Background

NR2C2AP may act as a repressor of NR2C2-mediated transactivation by suppressing the binding between NR2C2/TR4 and the TR4-response element in target genes.

NR2C2AP Antibody (N-term) - References

- Lamesch, P., et al. Genomics 89(3):307-315(2007)
Wang, C.P., et al. Biochem. Biophys. Res. Commun. 351(1):204-208(2006)
Lee, Y.F., et al. Biochem. Biophys. Res. Commun. 323(3):876-883(2004)
Yang, Y., et al. J. Biol. Chem. 278(9):7709-7717(2003)