

HIST1H2BL Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12710a**Specification**

HIST1H2BL Antibody (N-term) - Product Information

| | |
|-------------------|--|
| Application | FC, WB,E |
| Primary Accession | O99880 |
| Other Accession | O64524 , NP_003510.1 |
| Reactivity | Human |
| Predicted | Mouse |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 13952 |
| Antigen Region | 1-30 |

HIST1H2BL Antibody (N-term) - Additional Information**Gene ID** 8340**Other Names**

Histone H2B type 1-L, Histone H2Bc, H2B/c, HIST1H2BL, H2BFC

Target/Specificity

This HIST1H2BL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human HIST1H2BL.

Dilution

FC~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HIST1H2BL Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

HIST1H2BL Antibody (N-term) - Protein Information**Name** H2BC13 ([HGNC:4748](#))

Function Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location


Nucleus. Chromosome.

HIST1H2BL Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HIST1H2BL Antibody (N-term) - Images



HIST1H2BL Antibody (N-term) (Cat. #AP12710a) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the HIST1H2BL antibody detected the HIST1H2BL protein (arrow).



HIST1H2BL Antibody (N-term) (Cat. #AP12710a) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

HIST1H2BL Antibody (N-term) - Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H2B family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3.

HIST1H2BL Antibody (N-term) - References

Kim, S.C., et al. Mol. Cell 23(4):607-618(2006)
Beck, H.C., et al. Mol. Cell Proteomics 5(7):1314-1325(2006)
Pavri, R., et al. Cell 125(4):703-717(2006)
Bonenfant, D., et al. Mol. Cell Proteomics 5(3):541-552(2006)
Zhu, B., et al. Mol. Cell 20(4):601-611(2005)