

ZP4 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12724b

Specification

ZP4 Antibody (C-term) - Product Information

Application WB, IHC-P,E **Primary Accession** 012836 Other Accession NP 067009.1 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG **Antigen Region** 445-474

ZP4 Antibody (C-term) - Additional Information

Gene ID 57829

Other Names

Zona pellucida sperm-binding protein 4, Zona pellucida glycoprotein 4, Zp-4, Zona pellucida protein B, Processed zona pellucida sperm-binding protein 4, ZP4, ZPB

Target/Specificity

This ZP4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 445-474 amino acids from the C-terminal region of human ZP4.

Dilution

WB~~1:2000 IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ZP4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ZP4 Antibody (C-term) - Protein Information

Name ZP4

Synonyms ZPB





Tel: 858.875.1900 Fax: 858.875.1999

Function Component of the zona pellucida, an extracellular matrix surrounding oocytes which mediates sperm binding, induction of the acrosome reaction and prevents post-fertilization polyspermy. The zona pellucida is composed of 3 to 4 glycoproteins, ZP1, ZP2, ZP3, and ZP4. ZP4 may act as a sperm receptor.

Cellular Location

[Processed zona pellucida sperm-binding protein 4]: Zona pellucida {ECO:0000250|UniProtKB:Q00193}

Tissue Location


Expressed in oocytes.

ZP4 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ZP4 Antibody (C-term) - Images




ZP4 Antibody (C-term) (Cat. #AP12724b) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the ZP4 antibody detected the ZP4 protein (arrow).





Anti-ZP4 Antibody (C-term) at 1:2000 dilution + NCCIT whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 59 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



ZP4 Antibdy (C-term) (Cat. #AP12724b)immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of ZP4 Antibdy (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

ZP4 Antibody (C-term) - Background

The zona pellucida is an extracellular matrix that surrounds the oocyte and early embryo. It is composed primarily of three or four glycoproteins with various functions during fertilization and preimplantation development. The nascent protein contains a N-terminal signal peptide sequence, a conserved ZP domain, a consensus furin cleavage site, and a C-terminal transmembrane domain. It is hypothesized that furin cleavage results in release of the mature protein from the plasma membrane for subsequent incorporation into the zona pellucida matrix. However, the requirement for furin cleavage in this process remains controversial based on mouse studies. Previously, this gene has





Tel: 858.875.1900 Fax: 858.875.1999

been referred to as ZP1 or ZPB and thought to have similar functions as mouse Zp1. However, a human gene with higher similarity and chromosomal synteny to mouse Zp1 has been assigned the symbol ZP1 and this gene has been assigned the symbol ZP4.

ZP4 Antibody (C-term) - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010): McCauley, J.L., et al. Genes Immun. 10(7):624-630(2009) Nakano, M., et al. Proc. Natl. Acad. Sci. U.S.A. 106(31):12838-12842(2009) Choudhury, S., et al. J. Reprod. Immunol. 79(2):137-147(2009) Chiu, P.C., et al. Biol. Reprod. 79(5):869-877(2008) **ZP4** Antibody (C-term) - Citations

• Serum antibody immunoreactivity to equine zona protein after SpayVac vaccination.