

C6orf168 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12790b**Specification**

C6orf168 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q5TGI0
Other Accession	D3ZAT9 , Q3UMF9 , NP_115900.1
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	46843
Antigen Region	352-381

C6orf168 Antibody (C-term) - Additional Information**Gene ID** 84553**Other Names**

Failed axon connections homolog, FAXC, C6orf168

Target/Specificity

This C6orf168 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 352-381 amino acids from the C-terminal region of human C6orf168.

Dilution

WB~~1:1000

IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

C6orf168 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

C6orf168 Antibody (C-term) - Protein Information**Name** FAXC

Synonyms C6orf168

Function May play a role in axonal development.

Cellular Location


Membrane; Single-pass membrane protein

C6orf168 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C6orf168 Antibody (C-term) - Images



C6orf168 Antibody (C-term) (Cat. #AP12790b) western blot analysis in MCF-7 cell line lysates (35ug/lane). This demonstrates the C6orf168 antibody detected the C6orf168 protein (arrow).



C6orf168 Antibody (C-term) (Cat. #AP12790b) immunohistochemistry analysis in formalin fixed and paraffin embedded human testis tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of C6orf168 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

C6orf168 Antibody (C-term) - References

Stein, J.L., et al. Neuroimage 53(3):1160-1174(2010)
Mungall, A.J., et al. Nature 425(6960):805-811(2003)