

FCAR Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12915c**Specification**

FCAR Antibody (Center) - Product Information

Application	WB,E
Primary Accession	P24071
Other Accession	NP_579805.1 , NP_579803.1 , NP_001991.1 , NP_579813.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	32265
Antigen Region	58-84

FCAR Antibody (Center) - Additional Information**Gene ID** 2204**Other Names**

Immunoglobulin alpha Fc receptor, IgA Fc receptor, CD89, FCAR, CD89

Target/Specificity

This FCAR antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 58-84 amino acids from the Central region of human FCAR.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FCAR Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

FCAR Antibody (Center) - Protein Information**Name** FCAR

Synonyms CD89

Function Binds to the Fc region of immunoglobulins alpha. Mediates several functions including cytokine production.

Cellular Location

[Isoform A.1]: Cell membrane; Single-pass type I membrane protein [Isoform A.3]: Cell membrane; Single-pass type I membrane protein [Isoform B-delta-S2]: Secreted.

Tissue Location


Isoform A.1, isoform A.2 and isoform A.3 are differentially expressed between blood and mucosal myeloid cells Isoform A.1, isoform A.2 and isoform A.3 are expressed in monocytes Isoform A.1 and isoform A.2 are expressed in alveolar macrophages; however only one isoform is expressed at alveolar macrophages surfaces

FCAR Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FCAR Antibody (Center) - Images



Western blot analysis of FCAR (arrow) using rabbit polyclonal FCAR Antibody (Center) (Cat. #AP12915c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the FCAR gene.

FCAR Antibody (Center) - Background

This gene is a member of the immunoglobulin gene

superfamily and encodes a receptor for the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsonized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene.

FCAR Antibody (Center) - References

Vuong, M.T., et al. Kidney Int. (2010) In press :
Davila, S., et al. Genes Immun. 11(3):232-238(2010)
Peng, M., et al. Cell Res. 20(2):223-237(2010)
Kobayashi, T., et al. J. Dent. Res. 88(12):1137-1141(2009)
van der Steen, L., et al. Gastroenterology 137(6):2018-2029(2009)