

ELAVL2 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12937c

Specification

ELAVL2 Antibody (Center) - Product Information

IHC-P, FC, WB,E Application

Primary Accession 012926

Other Accession Q7SZT7, Q09032, Q61701, P26378, Q91584, Q60900, Q14576, Q91903, Q8CH84, Q60899,

NP 001164666.1, NP 001164668.1,

A0A0R4IEW8 Human, Mouse

Reactivity Predicted Rat, Xenopus, Zebrafish

Host **Rabbit** Clonality **Polyclonal** Rabbit IgG Isotype Antigen Region 156-184

ELAVL2 Antibody (Center) - Additional Information

Gene ID 1993

Other Names

ELAV-like protein 2, ELAV-like neuronal protein 1, Hu-antigen B, HuB, Nervous system-specific RNA-binding protein Hel-N1, ELAVL2, HUB

Target/Specificity

This ELAVL2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 156-184 amino acids from the Central region of human ELAVL2.

Dilution

IHC-P~~1:100 FC~~1:10~50 WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ELAVL2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.



ELAVL2 Antibody (Center) - Protein Information

Name ELAVL2

Synonyms HUB

Function RNA-binding protein that binds to the 3' untranslated region (3'UTR) of target mRNAs (By similarity). Seems to recognize a GAAA motif (By similarity). Can bind to its own 3'UTR, the FOS 3'UTR and the ID 3'UTR (By similarity).


Tissue LocationBrain; neural-specific.

ELAVL2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ELAVL2 Antibody (Center) - Images




ELAVL2 Antibody (Center) (Cat. #AP12937c) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the ELAVL2 antibody detected the ELAVL2 protein (arrow).






Anti-ELAVL2 Antibody (Center) at 1:1000 dilution + human brain lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 40 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Anti-ELAVL2 Antibody (Center) at 1:2000 dilution + Mouse brain lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 40 kDa Blocking/Dilution buffer: 5% NFDM/TBST.






Immunohistochemical analysis of AP12937C on paraffin-embedded Human brain tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of AP12937C on paraffin-embedded Human cerebellum tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.





ELAVL2 Antibody (Center) (Cat. #AP12937c) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ELAVL2 Antibody (Center) - Background

The protein encoded by this gene is a neural-specific RNA-binding protein that is known to bind to several 3' UTRs, including its own and also that of FOS and ID. The encoded protein may recognize a GAAA motif in the RNA. Three transcript variants encoding two different isoforms have been found for this gene.

ELAVL2 Antibody (Center) - References

Marroni, F., et al. Circ Cardiovasc Genet 2(4):322-328(2009) Lowe, J.K., et al. PLoS Genet. 5 (2), E1000365 (2009) : D'Alessandro, V., et al. Cell. Oncol. 30(4):291-297(2008) Jonson, L., et al. Mol. Cell Proteomics 6(5):798-811(2007) Yano, M., et al. J. Biol. Chem. 280(13):12690-12699(2005)