

DNAJC9 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12938A

Specification

DNAJC9 Antibody (N-term) - Product Information

WB, FC, E Application **Primary Accession 08WXX5** Other Accession NP 056005.1 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 29910 Antigen Region 29-58

DNAJC9 Antibody (N-term) - Additional Information

Gene ID 23234

Other Names

DnaJ homolog subfamily C member 9, DnaJ protein SB73, DNAJC9

Target/Specificity

This DNAJC9 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 29-58 amino acids from the N-terminal region of human DNAJC9.

Dilution

WB~~1:1000 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DNAJC9 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DNAJC9 Antibody (N-term) - Protein Information

Name DNAJC9

Function Acts as a dual histone chaperone and heat shock co-chaperone (PubMed: 33857403). As



a histone chaperone, forms a co-chaperone complex with MCM2 and histone H3-H4 heterodimers; and may thereby assist MCM2 in histone H3-H4 heterodimer recognition and facilitate the assembly of histones into nucleosomes (PubMed:33857403). May also act as a histone co-chaperone together with TONSL (PubMed:33857403). May recruit histone chaperones ASF1A, NASP and SPT2 to histone H3-H4 heterodimers (PubMed:33857403). Also plays a role as co-chaperone of the HSP70 family of molecular chaperone proteins, such as HSPA1A, HSPA1B and HSPA8 (PubMed:17182002, PubMed:33857403). As a co-chaperone, may play a role in the recruitment of HSP70-type molecular chaperone machinery to histone H3-H4 substrates, thereby maintaining the histone structural integrity (PubMed:33857403). Exhibits activity to assemble histones onto DNA in vitro (PubMed:33857403).

Cellular Location

Nucleus. Cytoplasm. Cell membrane. Note=Predominantly nuclear. Translocates to the cytoplasm and membrane after heat shock

Tissue Location


Expressed in heart, placenta, liver, skeletal muscle, kidney, pancreas, thymus, ovary, colon and peripheral blood

DNAJC9 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

DNAJC9 Antibody (N-term) - Images



DNAJC9 Antibody (N-term) (Cat. #AP12938a) western blot analysis in Jurkat cell line lysates (35ug/lane). This demonstrates the DNAJC9 antibody detected the DNAJC9 protein (arrow).





DNAJC9 Antibody (N-term) (Cat. #AP12938a) flow cytometric analysis of Jurkat cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated donkey-anti-rabbit secondary antibodies were used for the analysis.

DNAJC9 Antibody (N-term) - Background

DNAJC9 is upregulated at both mRNA and protein levels after various mitogenic and stress stimulations. DNAJC9 is mainly localized in cell nuclei under normal culture conditions, and transported into cytoplasm and plasma membrane upon heat shock stress through a non-classical and lipid-dependent pathway. DNAJC9 can interact with HSP70s and activate the ATPase activity of HSP70s, both of which are dependent on the J domain. Research has indicated that DNAJC9 is a novel cochaperone for HSP70s.

DNAJC9 Antibody (N-term) - References

Rose, J. Phd, et al. Mol. Med. (2010) In press:

Pan, Z., et al. Neoplasma 57(2):123-128(2010)

Han, C., et al. Biochem. Biophys. Res. Commun. 353(2):280-285(2007)

Takamura, Y., et al. Biochem. Biophys. Res. Commun. 285(2):387-392(2001)