

RASA1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12950b

Specification

RASA1 Antibody (C-term) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype Calculated MW Antigen Region WB,E <u>P20936</u> <u>P50904</u>, <u>NP_002881.1</u>, <u>NP_072179.1</u> Human Rat Rabbit Polyclonal Rabbit IgG 116403 966-993

RASA1 Antibody (C-term) - Additional Information

Gene ID 5921

Other Names Ras GTPase-activating protein 1, GAP, GTPase-activating protein, RasGAP, Ras p21 protein activator, p120GAP, RASA1, GAP, RASA

Target/Specificity

This RASA1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 966-993 amino acids from the C-terminal region of human RASA1.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RASA1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RASA1 Antibody (C-term) - Protein Information

Name RASA1



Synonyms GAP, RASA

Function Inhibitory regulator of the Ras-cyclic AMP pathway. Stimulates the GTPase of normal but not oncogenic Ras p21; this stimulation may be further increased in the presence of NCK1.

Cellular Location Cytoplasm.

Tissue Location

In placental villi, detected only in the trophoblast layer (cytotrophoblast and syncytiotrophoblast). Not detected in stromal, endothelial or Hofbauer cells (at protein level)

RASA1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

RASA1 Antibody (C-term) - Images



RASA1 Antibody (C-term) (Cat. #AP12950b) western blot analysis in 293 cell line lysates (35ug/lane).This demonstrates the RASA1 antibody detected the RASA1 protein (arrow).

RASA1 Antibody (C-term) - Background

The protein encoded by this gene is located in the cytoplasm and is part of the GAP1 family of GTPase-activating proteins. The gene product stimulates the GTPase activity of normal RAS p21 but not its oncogenic counterpart. Acting as a suppressor of RAS function, the protein enhances the weak intrinsic GTPase activity of RAS proteins resulting in the inactive GDP-bound form of RAS, thereby allowing control of cellular proliferation and differentiation. Mutations leading to changes in the binding sites



of either protein are associated with basal cell carcinomas. Alternative splicing results in two isoforms where the shorter isoform, lacking the N-terminal hydrophobic region but retaining the same activity, appears to be abundantly expressed in placental but not adult tissues.

RASA1 Antibody (C-term) - References

Hemerly, J.P., et al. Eur. J. Endocrinol. 163(5):747-755(2010) Wiemels, J.L., et al. Blood Cells Mol. Dis. 45(3):186-191(2010) Bachas, C., et al. Blood 116(15):2752-2758(2010) Oinuma, I., et al. J. Biol. Chem. 285(36):28200-28209(2010) Anand, S., et al. Nat. Med. 16(8):909-914(2010)