

PON2 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12952a

Specification

PON2 Antibody (N-term) - Product Information

Application	FC, IHC-P, WB,E
Primary Accession	Q15165
Other Accession	NP_000296.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	39381
Antigen Region	79-107

PON2 Antibody (N-term) - Additional Information

Gene ID 5445

Other Names

Serum paraoxonase/arylesterase 2, PON 2, Aromatic esterase 2, A-esterase 2, Serum arylalkylphosphatase 2, PON2

Target/Specificity

This PON2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 79-107 amino acids from the N-terminal region of human PON2.

Dilution

FC~~1:10~50

IHC-P~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PON2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PON2 Antibody (N-term) - Protein Information

Name PON2

Function Capable of hydrolyzing lactones and a number of aromatic carboxylic acid esters. Has antioxidant activity. Is not associated with high density lipoprotein. Prevents LDL lipid peroxidation, reverses the oxidation of mildly oxidized LDL, and inhibits the ability of MM-LDL to induce monocyte chemotaxis.

Cellular Location

Membrane; Peripheral membrane protein


Tissue Location

Widely expressed with highest expression in liver, lung, placenta, testis and heart.


PON2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PON2 Antibody (N-term) - Images

PON2 Antibody (N-term) (Cat. #AP12952a) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the PON2 antibody detected the PON2 protein (arrow).



PON2 Antibody (N-term) (Cat. #AP12952a) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of PON2 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



PON2 Antibody (N-term) (Cat. #AP12952a) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated donkey-anti-rabbit secondary antibodies were used for the analysis.

PON2 Antibody (N-term) - Background

This gene encodes a member of the paraoxonase gene family, which includes three known members located adjacent to each other on the long arm of chromosome 7. The encoded protein is ubiquitously expressed in human tissues, membrane-bound, and may act as a cellular antioxidant, protecting cells from oxidative stress. Hydrolytic activity against acylhomoserine lactones, important bacterial quorum-sensing mediators, suggests the encoded protein may also play a role in defense responses to pathogenic bacteria. Mutations in this gene may be associated with vascular disease and a number of quantitative phenotypes related to diabetes. Alternatively spliced transcript variants encoding different isoforms have been described.

PON2 Antibody (N-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Altenhofer, S., et al. J. Biol. Chem. 285(32):24398-24403(2010)
Ticozzi, N., et al. Ann. Neurol. 68(1):102-107(2010)
Wang, Y., et al. Diabet. Med. 27(4):376-383(2010)
Cross, D.S., et al. BMC Genet. 11, 51 (2010) :