

MRAP2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13187c**Specification**

MRAP2 Antibody (Center) - Product Information

| | |
|-------------------|-----------------------------|
| Application | IHC-P, WB,E |
| Primary Accession | O96G30 |
| Other Accession | NP_612418.2 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 23548 |
| Antigen Region | 58-86 |

MRAP2 Antibody (Center) - Additional Information**Gene ID** 112609**Other Names**

Melanocortin-2 receptor accessory protein 2, MC2R accessory protein 2, MRAP2, C6orf117

Target/Specificity

This MRAP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 58-86 amino acids from the Central region of human MRAP2.

Dilution

IHC-P~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MRAP2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

MRAP2 Antibody (Center) - Protein Information**Name** MRAP2

Synonyms C6orf117

Function Modulator of melanocortin receptor 4 (MC4R), a receptor involved in energy homeostasis. Plays a central role in the control of energy homeostasis and body weight regulation by increasing ligand- sensitivity of MC4R and MC4R-mediated generation of cAMP (By similarity). May also act as a negative regulator of MC2R: competes with MRAP for binding to MC2R and impairs the binding of corticotropin (ACTH) to MC2R. May also regulate activity of other melanocortin receptors (MC1R, MC3R and MC5R); however, additional evidence is required in vivo.

Cellular Location

Cell membrane; Single-pass membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein. Note=The formation of antiparallel homo- and heterodimers suggest that N- and C-terminus can both localize in the cytoplasmic and extracellular parts, depending on the context.

Tissue Location


Expressed in the adrenal gland and brain. Not expressed in other tissues.

MRAP2 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MRAP2 Antibody (Center) - Images



MRAP2 Antibody (Center) (Cat. #AP13187c) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the MRAP2 antibody detected the MRAP2 protein (arrow).



MRAP2 Antibody (Center) (Cat. #AP13187c) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of MRAP2 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

MRAP2 Antibody (Center) - Background

The specific function of this protein remains unknown.

MRAP2 Antibody (Center) - References

- Sebag, J.A., et al. Sci Signal 3 (116), RA28 (2010) :
- Chan, L.F., et al. Proc. Natl. Acad. Sci. U.S.A. 106(15):6146-6151(2009)
- Sebag, J.A., et al. J. Biol. Chem. 284(1):610-618(2009)
- Lamesch, P., et al. Genomics 89(3):307-315(2007)
- Metherell, L.A., et al. Nat. Genet. 37(2):166-170(2005)