

**OR2V2 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP13282b****Specification**

---

**OR2V2 Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O96R30</a>
Other Accession	<a href="#">NP_996763.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35339
Antigen Region	245-273

**OR2V2 Antibody (C-term) - Additional Information****Gene ID** 285659**Other Names**

Olfactory receptor 2V2, Olfactory receptor 2V3, Olfactory receptor OR5-3, OR2V2, OR2V3

**Target/Specificity**

This OR2V2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 245-273 amino acids from the C-terminal region of human OR2V2.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

OR2V2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**OR2V2 Antibody (C-term) - Protein Information****Name** OR2V2**Synonyms** OR2V3

**Function** Odorant receptor.

**Cellular Location**


Cell membrane; Multi-pass membrane protein.

**OR2V2 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**OR2V2 Antibody (C-term) - Images**



OR2V2 Antibody (C-term) (Cat. #AP13282b) western blot analysis in human normal Uterus tissue lysates (35ug/lane). This demonstrates the OR2V2 antibody detected the OR2V2 protein (arrow).

**OR2V2 Antibody (C-term) - Background**

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

**OR2V2 Antibody (C-term) - References**

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004) Fuchs, T., et al. Genomics 80(3):295-302(2002)