

HtrA3 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1332b**Specification**

HtrA3 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P83110
Other Accession	D3ZA76
Reactivity	Human
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	48608
Antigen Region	394-425

HtrA3 Antibody (C-term) - Additional Information**Gene ID** 94031**Other Names**

Serine protease HTRA3, 3421-, High-temperature requirement factor A3, Pregnancy-related serine protease, HTRA3, PRSP

Target/Specificity

This HtrA3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 394-425 amino acids from the C-terminal region of human HtrA3.

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HtrA3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

HtrA3 Antibody (C-term) - Protein Information**Name** HTRA3

Synonyms PRSP

Function Serine protease that cleaves beta-casein/CSN2 as well as several extracellular matrix (ECM) proteoglycans such as decorin/DCN, biglycan/BGN and fibronectin/FN1. Inhibits signaling mediated by TGF- beta family proteins possibly indirectly by degradation of these ECM proteoglycans (By similarity). May act as a tumor suppressor. Negatively regulates, in vitro, trophoblast invasion during placental development and may be involved in the development of the placenta in vivo. May also have a role in ovarian development, granulosa cell differentiation and luteinization (PubMed:[21321049](#), PubMed:[2229724](#)).

Cellular Location

Secreted. Note=Secretion increased during decidualization of endometrial stromal cells

Tissue Location


Widely expressed, with highest levels in both adult and fetal heart, ovary, uterus placenta, and bladder. In the endometrium, expressed in epithelial glands and the stroma. Also present in leukocytes. Isoform 1 is predominant in heart and skeletal muscle, whereas isoform 2 is predominant in placenta and kidney

HtrA3 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HtrA3 Antibody (C-term) - Images



Western blot analysis of anti-HtrA3 C-term Pab (Cat. #AP1332b) in mouse brain tissue lysate. HtrA3 (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

HtrA3 Antibody (C-term) - Background

Insulin-like growth factors (IGFs) stimulate the proliferation and differentiation of a vast number of cell types. The action of the growth factors is mediated and controlled by a complex system of components, including several proteases that cleave the IGF-Binding Proteins. HtrA1 is a 480 aa protein that contains an N-terminus homologous to MAC25 (IGFBP7) with a conserved Kazal-type serine protease inhibitor motif, as well as a C-terminal PDZ domain. Semiquantitative RT-PCR and immunoblot analyses showed an approximately 7-fold increase of PRSS11 in osteoarthritis cartilage compared with controls. HTRA2 is released from mitochondria and inhibits the function of XIAP by direct binding in a way similar to SMAC. Moreover, when overexpressed extramitochondrially, HTRA2 induced atypical cell death, which was neither accompanied by a significant increase in caspase activity nor inhibited by caspase inhibitors, including XIAP. A catalytically inactive mutant of HTRA2, however, did not induce cell death. Suzuki et al. (2001) concluded that HTRA2 is a SMAC-like inhibitor of IAP (inhibitor of apoptosis proteins) activity with a serine protease-dependent cell death-inducing activity.

HtrA3 Antibody (C-term) - References

Nie, G.Y., et al., Biochem. J. 371 (Pt 1), 39-48 (2003).