

UGT2B11 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13518b

Specification

UGT2B11 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O75310
Other Accession	O97951 , P36537 , O02663 , P16662 , NP_001064.1
Reactivity	Human
Predicted	Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	61038
Antigen Region	429-458

UGT2B11 Antibody (C-term) - Additional Information

Gene ID 10720

Other Names

UDP-glucuronosyltransferase 2B11, UDPGT 2B11, UGT2B11

Target/Specificity

This UGT2B11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 429-458 amino acids from the C-terminal region of human UGT2B11.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

UGT2B11 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

UGT2B11 Antibody (C-term) - Protein Information

Name UGT2B11

Function UDPGT is of major importance in the conjugation and subsequent elimination of potentially toxic xenobiotics and endogenous compounds.

Cellular Location

Microsome membrane; Single-pass membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein

Tissue Location


Widely expressed.

UGT2B11 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UGT2B11 Antibody (C-term) - Images



UGT2B11 Antibody (C-term) (Cat. #AP13518b) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the UGT2B11 antibody detected the UGT2B11 protein (arrow).

UGT2B11 Antibody (C-term) - Background

UDPGT is of major importance in the conjugation and subsequent elimination of potentially toxic xenobiotics and endogenous compounds.

UGT2B11 Antibody (C-term) - References

Li, J., et al. Breast Cancer Res. 12 (2), R19 (2010) :
Ross, C.J., et al. Nat. Genet. 41(12):1345-1349(2009)
Saito, A., et al. J. Hum. Genet. 54(6):317-323(2009)
Beaulieu, M., et al. Biochem. Biophys. Res. Commun. 248(1):44-50(1998)
Jin, C.J., et al. Biochem. Biophys. Res. Commun. 194(1):496-503(1993)