

UBN2 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13523a**Specification**

UBN2 Antibody (N-term) - Product Information

Application	WB, FC,E
Primary Accession	Q6ZU65
Other Accession	NP_775840.3
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	146089
Antigen Region	171-200

UBN2 Antibody (N-term) - Additional Information**Gene ID** 254048**Other Names**

Ubinuclein-2, UBN2, KIAA2030

Target/Specificity

This UBN2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 171-200 amino acids from the N-terminal region of human UBN2.

Dilution

WB~~1:1000

FC~~1:10~50

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

UBN2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

UBN2 Antibody (N-term) - Protein Information**Name** UBN2

Synonyms KIAA2030

Tissue Location


Expressed in several cell lines tested, including primary and transformed cell lines.

UBN2 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


UBN2 Antibody (N-term) - Images



UBN2 Antibody (N-term) (Cat. #AP13523a) western blot analysis in mouse stomach tissue lysates (35ug/lane). This demonstrates the UBN2 antibody detected the UBN2 protein (arrow).



All lanes : Anti-UBN2 Antibody (N-term) at 1:1000 dilution Lane 1: Ramos whole cell lysate Lane 2: U-2OS whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 146 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



UBN2 Antibody (N-term) (Cat. #AP13523a) flow cytometric analysis of Ramos cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated donkey-anti-rabbit secondary antibodies were used for the analysis.

UBN2 Antibody (N-term) - Background

The specific function of the protein remains unknown.

UBN2 Antibody (N-term) - References

Banumathy, G., et al. Mol. Cell. Biol. 29(3):758-770(2009)