

**TUBA4A Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP13535b**

**Specification**

---

**TUBA4A Antibody (C-term) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">P68366</a>
Other Accession	<a href="#">Q5XIF6</a> , <a href="#">P68368</a> , <a href="#">P68367</a> , <a href="#">P81948</a> , <a href="#">NP_005991.1</a>
Reactivity	Human, Mouse, Rat
Predicted	Bovine, Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	417-446

**TUBA4A Antibody (C-term) - Additional Information**

**Gene ID** 7277

**Other Names**

Tubulin alpha-4A chain, Alpha-tubulin 1, Testis-specific alpha-tubulin, Tubulin H2-alpha, Tubulin alpha-1 chain, TUBA4A, TUBA1

**Target/Specificity**

This TUBA4A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 417-446 amino acids from the C-terminal region of human TUBA4A.

**Dilution**

WB~~1:8000  
IHC-P~~1:25

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

TUBA4A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**TUBA4A Antibody (C-term) - Protein Information**

**Name** TUBA4A

## Synonyms TUBA1

**Function** Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha chain.

## Cellular Location


Cytoplasm, cytoskeleton.

## TUBA4A Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## TUBA4A Antibody (C-term) - Images



All lanes : Anti-TUBA4A Antibody (C-term) at 1:8000 dilution Lane 1: A431 whole cell lysate Lane 2: HeLa whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: MCF-7 whole cell lysate Lane 5: NIH/3T3 whole cell lysate Lane 6: Mouse brain lysate Lane 7: PC-12 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 50 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



AP13535B staining TUBA4A in human lung tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

#### **TUBA4A Antibody (C-term) - Background**

Microtubules of the eukaryotic cytoskeleton perform essential and diverse functions and are composed of a heterodimer of alpha and beta tubulin. The genes encoding these microtubule constituents are part of the tubulin superfamily, which is composed of six distinct families. Genes from the alpha, beta and gamma tubulin families are found in all eukaryotes. The alpha and beta tubulins represent the major components of microtubules, while gamma tubulin plays a critical role in the nucleation of microtubule assembly. There are multiple alpha and beta tubulin genes and they are highly conserved among and between species. This gene encodes an alpha tubulin that is a highly conserved homolog of a rat testis-specific alpha tubulin.

#### **TUBA4A Antibody (C-term) - References**

Houck, S.A., et al. PLoS ONE 5 (7), E11795 (2010) :  
Zhao, J., et al. BMC Med. Genet. 11, 96 (2010) :  
Gudbjartsson, D.F., et al. Nat. Genet. 40(5):609-615(2008)  
Petretti, C., et al. EMBO Rep. 7(4):418-424(2006)  
Fiore, G., et al. Neurosci. Lett. 394(1):57-62(2006)

#### **TUBA4A Antibody (C-term) - Citations**

- [Dysregulation of a novel miR-1825/TBCB/TUBA4A pathway in sporadic and familial ALS.](#)