

APOBEC3B Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1353a**Specification**

APOBEC3B Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O9UH17
Other Accession	P31941
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	299-330

APOBEC3B Antibody (C-term) - Additional Information**Gene ID** 9582**Other Names**

DNA dC->dU-editing enzyme APOBEC-3B, A3B, 354-, Phorbolin-1-related protein, Phorbolin-2/3, APOBEC3B

Target/Specificity

This APOBEC3B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 299-330 amino acids from the C-terminal region of human APOBEC3B.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

APOBEC3B Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

APOBEC3B Antibody (C-term) - Protein Information**Name** APOBEC3B**Function** DNA deaminase (cytidine deaminase) which acts as an inhibitor of retrovirus replication

and retrotransposon mobility via deaminase- dependent and -independent mechanisms. After the penetration of retroviral nucleocapsids into target cells of infection and the initiation of reverse transcription, it can induce the conversion of cytosine to uracil in the minus-sense single-strand viral DNA, leading to G-to-A hypermutations in the subsequent plus-strand viral DNA. The resultant detrimental levels of mutations in the proviral genome, along with a deamination-independent mechanism that works prior to the proviral integration, together exert efficient antiretroviral effects in infected target cells. Selectively targets single-stranded DNA and does not deaminate double-stranded DNA or single- or double-stranded RNA. Exhibits antiviral activity against simian immunodeficiency virus (SIV), hepatitis B virus (HBV) and human T-cell leukemia virus type 1 (HTLV-1) and may inhibit the mobility of LTR and non-LTR retrotransposons.

Cellular Location

Nucleus

Tissue Location


Expressed at high and moderate levels in peripheral blood leukocytes, spleen, testes, heart, thymus, prostate and ovary Also expressed at low levels in several other tissues

APOBEC3B Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

APOBEC3B Antibody (C-term) - Images



Western blot analysis of hAPOBEC3B(Cat. #AP1353a) in mouse testis tissue lysates (35ug/lane). APOBEC3B (arrow) was detected using the purified Pab.

APOBEC3B Antibody (C-term) - Citations

- [APOBEC3B: A Potential Factor Suppressing Growth of Human Hepatocellular Carcinoma Cells.](#)

