

### MMP28 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP13734c

## **Specification**

### MMP28 Antibody (Center) - Product Information

Application IHC-P, WB,E Primary Accession Q9H239

Other Accession NP 001027449.1, NP 077278.1

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Human
Rabbit
Polyclonal
Rabbit IgG
195-223

## MMP28 Antibody (Center) - Additional Information

#### **Gene ID** 79148

#### **Other Names**

Matrix metalloproteinase-28, MMP-28, 3424-, Epilysin, MMP28, MMP25

#### Target/Specificity

This MMP28 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 195-223 amino acids from the Central region of human MMP28.

# **Dilution**

IHC-P~~1:10~50 WB~~1:1000

E~~Use at an assay dependent concentration.

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

MMP28 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

# MMP28 Antibody (Center) - Protein Information

#### Name MMP28



# **Synonyms MMP25**

Function Can degrade casein. Could play a role in tissues homeostasis and repair.

#### **Cellular Location**

Secreted, extracellular space, extracellular matrix

#### **Tissue Location**


Expressed at high levels in testes and lung. Low levels are detected in kidney, pancreas and skin. Also expressed in fetal lung, brain, skeletal muscle and kidney. Expressed selectively in keratinocytes. Widely expressed in several carcinomas as well. Is up- regulated in response to injury in the skin

## MMP28 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# MMP28 Antibody (Center) - Images



MMP28 Antibody (Center) (Cat. #AP13734c) western blot analysis in MDA-MB435 cell line lysates (35ug/lane). This demonstrates the MMP28 antibody detected the MMP28 protein (arrow).





MMP28 Antibody (Center) (Cat. #AP13734c)immunohistochemistry analysis in formalin fixed and paraffin embedded human pancreas tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of MMP28 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

# MMP28 Antibody (Center) - Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix for both normal physiological processes, such as embryonic development, reproduction and tissue remodeling, and disease processes, such as asthma and metastasis. This gene encodes a secreted enzyme that degrades casein. Its expression pattern suggests that it plays a role in tissue homeostasis and in wound repair. Transcript variants encoding different isoforms have been described. [provided by RefSeq].

# MMP28 Antibody (Center) - References

Swingler, T.E., et al. Biochem. J. 427(3):391-400(2010) Suomela, S., et al. Virchows Arch. 455(6):495-503(2009) Rodgers, U.R., et al. Matrix Biol. 28(5):263-272(2009) Heiskanen, T.J., et al. Matrix Biol. 28(2):74-83(2009) Gruber, H.E., et al. Arthritis Res. Ther. 11 (6), R184 (2009) :