

**CUL4A Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP13873a**

**Specification**

---

**CUL4A Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q13619</a>
Other Accession	<a href="#">Q3TCH7</a> , <a href="#">NP_001008895.1</a> , <a href="#">NP_003580.1</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	87680
Antigen Region	31-60

**CUL4A Antibody (N-term) - Additional Information**

**Gene ID** 8451

**Other Names**

Cullin-4A, CUL-4A, CUL4A

**Target/Specificity**

This CUL4A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 31-60 amino acids from the N-terminal region of human CUL4A.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CUL4A Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CUL4A Antibody (N-term) - Protein Information**

**Name** CUL4A {ECO:0000303|PubMed:9721878, ECO:0000312|HGNC:HGNC:2554}

**Function** Core component of multiple cullin-RING-based E3 ubiquitin- protein ligase complexes


which mediate the ubiquitination of target proteins (PubMed:[14578910](#), PubMed:[14739464](#), PubMed:[15448697](#), PubMed:[15548678](#), PubMed:[15811626](#), PubMed:[16678110](#), PubMed:[17041588](#), PubMed:[24209620](#), PubMed:[30166453](#), PubMed:[33854232](#), PubMed:[33854239](#)). As a scaffold protein may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme (PubMed:[14578910](#), PubMed:[14739464](#), PubMed:[15448697](#), PubMed:[15548678](#), PubMed:[15811626](#), PubMed:[16678110](#), PubMed:[17041588](#), PubMed:[24209620](#)). The E3 ubiquitin- protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1 (PubMed:[14578910](#), PubMed:[14739464](#), PubMed:[15448697](#), PubMed:[15548678](#), PubMed:[15811626](#), PubMed:[16678110](#), PubMed:[17041588](#), PubMed:[24209620](#)). The functional specificity of the E3 ubiquitin-protein ligase complex depends on the variable substrate recognition component (PubMed:[14578910](#), PubMed:[14739464](#), PubMed:[15448697](#), PubMed:[15548678](#), PubMed:[15811626](#), PubMed:[16678110](#), PubMed:[17041588](#), PubMed:[24209620](#)). DCX(DET1-COP1) directs ubiquitination of JUN (PubMed:[14739464](#)). DCX(DDB2) directs ubiquitination of XPC (PubMed:[15811626](#)). DCX(DDB2) ubiquitinates histones H3-H4 and is required for efficient histone deposition during replication-coupled (H3.1) and replication-independent (H3.3) nucleosome assembly, probably by facilitating the transfer of H3 from ASF1A/ASF1B to other chaperones involved in histone deposition (PubMed:[16678110](#), PubMed:[17041588](#), PubMed:[24209620](#)). DCX(DTL) plays a role in PCNA-dependent polyubiquitination of CDT1 and MDM2-dependent ubiquitination of p53/TP53 in response to radiation-induced DNA damage and during DNA replication (PubMed:[14578910](#), PubMed:[15448697](#), PubMed:[15548678](#)). DCX(DTL) directs autoubiquitination of DTL (PubMed:[23478445](#)). In association with DDB1 and SKP2 probably is involved in ubiquitination of CDKN1B/p27kip (PubMed:[16537899](#)). Is involved in ubiquitination of HOXA9 (PubMed:[14609952](#)). The DDB1-CUL4A- DTL E3 ligase complex regulates the circadian clock function by mediating the ubiquitination and degradation of CRY1 (PubMed:[26431207](#)). The DCX(ERCC8) complex (also named CSA complex) plays a role in transcription-coupled repair (TCR) (PubMed:[12732143](#), PubMed:[32355176](#), PubMed:[38316879](#)). A number of DCX complexes (containing either TRPC4AP or DCAF12 as substrate-recognition component) are part of the DesCEND (destruction via C-end degrons) pathway, which recognizes a C-degron located at the extreme C terminus of target proteins, leading to their ubiquitination and degradation (PubMed:[29779948](#)). The DCX(AMBRA1) complex is a master regulator of the transition from G1 to S cell phase by mediating ubiquitination of phosphorylated cyclin-D (CCND1, CCND2 and CCND3) (PubMed:[33854232](#), PubMed:[33854239](#)). The DCX(AMBRA1) complex also acts as a regulator of Cul5-RING (CRL5) E3 ubiquitin-protein ligase complexes by mediating ubiquitination and degradation of Elongin-C (ELOC) component of CRL5 complexes (PubMed:[30166453](#)). With CUL4B, contributes to ribosome biogenesis (PubMed:[26711351](#)).


## CUL4A Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## CUL4A Antibody (N-term) - Images






CUL4A Antibody (N-term) (Cat. #AP13873a) western blot analysis in mouse testis tissue lysates (35ug/lane). This demonstrates the CUL4A antibody detected the CUL4A protein (arrow).



CUL4A Antibody (N-term) (Cat. #AP13873a) western blot analysis in Jurkat cell line lysates (35ug/lane). This demonstrates the CUL4A antibody detected the CUL4A protein (arrow).



Western blot analysis of CUL4A (arrow) using rabbit polyclonal CUL4A Antibody (N-term) (Cat. #AP13873a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the CUL4A gene.

#### **CUL4A Antibody (N-term) - Background**

CUL4A is the ubiquitin ligase component of a multimeric complex involved in the degradation of DNA damage-response proteins

(Liu et al., 2009 [PubMed 19481525]).

#### **CUL4A Antibody (N-term) - References**

Aggarwal, P., et al. Cancer Cell 18(4):329-340(2010)

Abbas, T., et al. Mol. Cell 40(1):9-21(2010)

Lv, X.B., et al. J. Biol. Chem. 285(24):18234-18240(2010)

Kerzendorfer, C., et al. Hum. Mol. Genet. 19(7):1324-1334(2010)

Melchor, L., et al. Breast Cancer Res. 11 (6), R86 (2009) :