

**SLC30A1 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP13972b**

### Specification

#### SLC30A1 Antibody (C-term) - Product Information

|                   |  |
|-------------------|--|
| Application       | WB,E   |
| Primary Accession | <a href="#">Q9Y6M5</a>                               |
| Other Accession   | <a href="#">Q4R6K2</a> , <a href="#">NP_067017.2</a> |
| Reactivity        | Human  |
| Predicted         | Monkey   |
| Host              | Rabbit   |
| Clonality         | Polyclonal   |
| Isotype           | Rabbit IgG   |
| Antigen Region    | 458-487  |

#### SLC30A1 Antibody (C-term) - Additional Information

##### Gene ID 7779

##### Other Names

Zinc transporter 1, ZnT-1, Solute carrier family 30 member 1, SLC30A1, ZNT1

##### Target/Specificity

This SLC30A1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 458-487 amino acids from the C-terminal region of human SLC30A1.

##### Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

##### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

##### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions

SLC30A1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### SLC30A1 Antibody (C-term) - Protein Information

Name SLC30A1 ([HGNC:11012](#))

Function Zinc ion:proton antiporter that could function at the plasma membrane mediating zinc

efflux from cells against its electrochemical gradient protecting them from intracellular zinc accumulation and toxicity (PubMed:[31471319](#)). Alternatively, could prevent the transport to the plasma membrane of CACNB2, the L-type calcium channels regulatory subunit, through a yet to be defined mechanism. By modulating the expression of these channels at the plasma membrane, could prevent calcium and zinc influx into cells. By the same mechanism, could also prevent L-type calcium channels-mediated heavy metal influx into cells (By similarity). In some cells, could also function as a zinc ion:proton antiporter mediating zinc entry into the lumen of cytoplasmic vesicles. In macrophages, can increase zinc ions concentration into the lumen of cytoplasmic vesicles containing engulfed bacteria and could help inactivate them (PubMed:[32441444](#)). Forms a complex with TMC6/EVER1 and TMC8/EVER2 at the ER membrane of keratinocytes which facilitates zinc uptake into the ER (PubMed:[18158319](#)). Down-regulates the activity of transcription factors induced by zinc and cytokines (PubMed:[18158319](#)).

### **Cellular Location**


Cell membrane; Multi-pass membrane protein. Basolateral cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane; Multi-pass membrane protein. Cytoplasm. Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus membrane; Multi-pass membrane protein. Nucleus membrane; Multi- pass membrane protein. Note=Localization to the plasma membrane is regulated by cellular zinc status. Recruitment to the plasma membrane from an internal pool is stimulated by zinc while in absence of zinc the plasma membrane pool is endocytosed and degraded (PubMed:31471319). Localizes to the basolateral surface of enterocytes (By similarity). Localizes to zinc-containing intracellular vesicles in macrophages (PubMed:32441444). Localizes in the cytoplasm and to the ER, Golgi and nucleus membranes in keratinocytes (PubMed:18158319) {ECO:0000250|UniProtKB:Q62720, ECO:0000269|PubMed:18158319, ECO:0000269|PubMed:31471319, ECO:0000269|PubMed:32441444}

### **SLC30A1 Antibody (C-term) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **SLC30A1 Antibody (C-term) - Images**



SLC30A1 Antibody (C-term) (Cat. #AP13972b) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the SLC30A1 antibody detected the SLC30A1 protein (arrow).



Anti-SLC30A1 Antibody (C-term) at 1:1000 dilution + Hela whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 55 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

### SLC30A1 Antibody (C-term) - Background

SLC30A1 may be involved in zinc transport out of the cell.

### SLC30A1 Antibody (C-term) - References

Urani, C., et al. Toxicol In Vitro 24(2):370-374(2010)  
Beharier, O., et al. Ann. N. Y. Acad. Sci. 1188, 87-95 (2010) :  
Guey, L.T., et al. Eur. Urol. 57(2):283-292(2010)  
Hosgood, H.D. III, et al. Respir Med 103(12):1866-1870(2009)  
Levy, S., et al. J. Biol. Chem. 284(47):32434-32443(2009)

### SLC30A1 Antibody (C-term) - Citations

- [Zinc transporters ZnT3 and ZnT6 are downregulated in the spinal cords of patients with sporadic amyotrophic lateral sclerosis.](#)