

NOC3L Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13976a**Specification**

NOC3L Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q8WTT2
Other Accession	NP_071896.8
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	92548
Antigen Region	49-78

NOC3L Antibody (N-term) - Additional Information**Gene ID** 64318**Other Names**

Nucleolar complex protein 3 homolog, NOC3 protein homolog, Factor for adipocyte differentiation 24, NOC3-like protein, Nucleolar complex-associated protein 3-like protein, NOC3L, AD24, C10orf117, FAD24

Target/Specificity

This NOC3L antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 49-78 amino acids from the N-terminal region of human NOC3L.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NOC3L Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NOC3L Antibody (N-term) - Protein Information**Name** NOC3L

Synonyms AD24, C10orf117, FAD24

Function May be required for adipogenesis.

Cellular Location

Nucleus, nucleolus. Nucleus speckle.

Tissue Location


Expressed in colon, heart, kidney, liver, lung, placenta, skeletal muscle, small intestine, spleen and thymus

NOC3L Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NOC3L Antibody (N-term) - Images



Western blot analysis of lysate from SK-N-MC cell line, using NOC3L Antibody (N-term)(Cat. #AP13976a). AP13976a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug per lane.

NOC3L Antibody (N-term) - Background

NOC3L may be required for adipogenesis (By similarity).

NOC3L Antibody (N-term) - References

Nousiainen, M., et al. Proc. Natl. Acad. Sci. U.S.A. 103(14):5391-5396(2006)

Nousiainen, M., et al. Proc. Natl. Acad. Sci. U.S.A. 103(14):5391-5396(2006)

Grupe, A., et al. Am. J. Hum. Genet. 78(1):78-88(2006)

Tominaga, K., et al. J. Cell. Sci. 117 (PT 25), 6217-6226 (2004) :

Deloukas, P., et al. Nature 429(6990):375-381(2004)