

**TICAM2 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP14060a**

**Specification**

---

**TICAM2 Antibody (N-term) - Product Information**

|                   |  |
|-------------------|--|
| Application       | WB,E   |
| Primary Accession | <a href="#">Q86XR7</a>                                       |
| Other Accession   | <a href="#">NP_001157941.1</a> , <a href="#">NP_067681.1</a> |
| Reactivity        | Human  |
| Host              | Rabbit   |
| Clonality         | Polyclonal   |
| Isotype           | Rabbit IgG   |
| Calculated MW     | 26916  |
| Antigen Region    | 12-41  |

**TICAM2 Antibody (N-term) - Additional Information**

**Gene ID** 100302736;353376

**Other Names**

TIR domain-containing adapter molecule 2, TICAM-2, Putative NF-kappa-B-activating protein 502, TRIF-related adapter molecule, Toll-like receptor adaptor protein 3, Toll/interleukin-1 receptor domain-containing protein, MyD88-4, TICAM2, TIRAP3, TIRP, TRAM

**Target/Specificity**

This TICAM2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 12-41 amino acids from the N-terminal region of human TICAM2.

**Dilution**

WB~~1:1000  
E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

TICAM2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**TICAM2 Antibody (N-term) - Protein Information**

**Name** TICAM2

## Synonyms TIRAP3, TIRP, TRAM

**Function** Functions as a sorting adapter in different signaling pathways to facilitate downstream signaling leading to type I interferon induction (PubMed:[16603631](#), PubMed:[16757566](#), PubMed:[25385819](#), PubMed:[25825441](#)). In TLR4 signaling, physically bridges TLR4 and TICAM1 and functionally transmits signal to TICAM1 in early endosomes after endocytosis of TLR4. In TLR2 signaling, physically bridges TLR2 and MYD88 and is required for the TLR2- dependent movement of MYD88 to endosomes following ligand engagement (PubMed:[25385819](#)). Involved in IL-18 signaling and is proposed to function as a sorting adapter for MYD88 in IL-18 signaling during adaptive immune response (PubMed:[22685567](#)). Forms a complex with RAB11FIP2 that is recruited to the phagosomes to promote the activation of the actin-regulatory GTPases RAC1 and CDC42 and subsequent phagocytosis of Gram-negative bacteria (PubMed:[30883606](#)).

## Cellular Location

[Isoform 1]: Cytoplasm. Golgi apparatus. Cell membrane. Endoplasmic reticulum. Early endosome membrane. Late endosome membrane. Cell projection, phagocytic cup. Note=Localized to the plasma membrane as a result of myristoylation. Phosphorylation on Ser-16 leads to its depletion from the membrane. Upon LPS stimulation colocalizes with isoform 2 in late endosomes

## Tissue Location


Expressed in spleen, prostate, testis, uterus, small intestine, colon, peripheral blood leukocytes, heart, placenta, lung, liver, skeletal muscle, and pancreas Isoform 2 is ubiquitously expressed (at lower levels than isoform 1)


## TICAM2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## TICAM2 Antibody (N-term) - Images





TICAM2 Antibody (N-term) (Cat. #AP14060a) western blot analysis in human placenta tissue lysates (35ug/lane). This demonstrates the TICAM2 antibody detected the TICAM2 protein (arrow).

#### **TICAM2 Antibody (N-term) - Background**

TIRP is a Toll/interleukin-1 receptor (IL1R; MIM 147810) (TIR) domain-containing adaptor protein involved in Toll receptor signaling (see TLR4; MIM 603030).

#### **TICAM2 Antibody (N-term) - References**

Lysakova-Devine, T., et al. J. Immunol. 185(7):4261-4271(2010)  
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :  
Palsson-McDermott, E.M., et al. Nat. Immunol. 10(6):579-586(2009)  
Hawn, T.R., et al. PLoS ONE 4 (6), E5990 (2009) :  
Nakajima, T., et al. Immunogenetics 60(12):727-735(2008)