

Mouse Map2k7 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14155c**Specification**

Mouse Map2k7 Antibody (Center) - Product Information

Application	IHC-P, WB,E
Primary Accession	Q8CE90
Other Accession	Q4KSH7 , O14733 , NP_001157644.1 , NP_001036022.1
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	59312
Antigen Region	206-233

Mouse Map2k7 Antibody (Center) - Additional Information**Gene ID** 26400**Other Names**

Dual specificity mitogen-activated protein kinase kinase 7, MAP kinase kinase 7, MAPKK 7, JNK-activating kinase 2, MAPK/ERK kinase 7, MEK 7, c-Jun N-terminal kinase kinase 2, JNK kinase 2, JNKK 2, Map2k7 {ECO:0000312|MGI:MGI:1346871}

Target/Specificity

This Mouse Map2k7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 206-233 amino acids from the Central region of mouse Map2k7.

Dilution

IHC-P~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Map2k7 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Map2k7 Antibody (Center) - Protein Information

Name Map2k7 {ECO:0000312|MGI:MGI:1346871}

Function Dual specificity protein kinase which acts as an essential component of the MAP kinase signal transduction pathway. Essential component of the stress-activated protein kinase/c-Jun N-terminal kinase (SAP/JNK) signaling pathway. With MAP2K4/MKK4, is the one of the only known kinase to directly activate the stress-activated protein kinase/c-Jun N-terminal kinases MAPK8/JNK1, MAPK9/JNK2 and MAPK10/JNK3. MAP2K4/MKK4 and MAP2K7/MKK7 both activate the JNKs by phosphorylation, but they differ in their preference for the phosphorylation site in the Thr-Pro-Tyr motif. MAP2K4/MKK4 shows preference for phosphorylation of the Tyr residue and MAP2K7/MKK7 for the Thr residue. The monophosphorylation of JNKs on the Thr residue is sufficient to increase JNK activity indicating that MAP2K7/MKK7 is important to trigger JNK activity, while the additional phosphorylation of the Tyr residue by MAP2K4/MKK4 ensures optimal JNK activation. Has a specific role in JNK signal transduction pathway activated by pro-inflammatory cytokines. The MKK/JNK signaling pathway is also involved in mitochondrial death signaling pathway, including the release cytochrome c, leading to apoptosis. Part of a non-canonical MAPK signaling pathway, composed of the upstream MAP3K12 kinase and downstream MAP kinases MAPK1/ERK2 and MAPK3/ERK1, that enhances the AP-1-mediated transcription of APP in response to APOE (PubMed:[28111074](#)).

Cellular Location

Nucleus. Cytoplasm

Tissue Location

Expressed at high levels in brain, lung, liver, skeletal muscle, kidney, and testis and at lower levels in the heart and spleen.


Mouse Map2k7 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mouse Map2k7 Antibody (Center) - Images






Mouse Map2k7 Antibody (Center) (Cat. #AP14155c) western blot analysis in Jurkat cell line lysates (35ug/lane). This demonstrates the Map2k7 antibody detected the Map2k7 protein (arrow).



Mouse Map2k7 Antibody (Center) (Cat. #AP14155c) western blot analysis in mouse liver tissue lysates (35ug/lane). This demonstrates the Map2k7 antibody detected the Map2k7 protein (arrow).



Mouse Map2k7 Antibody (Center) (AP14155c) immunohistochemistry analysis in formalin fixed and paraffin embedded mouse brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of Mouse Map2k7 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

Mouse Map2k7 Antibody (Center) - Background

Stress activated, dual specificity kinase that activates the JUN kinases MAPK8/JNK1, MAPK9/JNK2 and MAPK10/JNK3.

Mouse Map2k7 Antibody (Center) - References

Bogani, D., et al. PLoS Biol. 7 (9), E1000196 (2009) :
Derradji, H., et al. Dev. Biol. 322(2):302-313(2008)
Nishitai, G., et al. J. Cell. Biochem. 104(5):1771-1780(2008)
Wang, X., et al. Mol. Cell. Biol. 27(22):7935-7946(2007)
Jaeschke, A., et al. Mol. Cell 27(3):498-508(2007)