

TACR3 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14222b**Specification**

TACR3 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P29371
Other Accession	NP_001050.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	52202
Antigen Region	406-435

TACR3 Antibody (C-term) - Additional Information**Gene ID** 6870**Other Names**

Neuromedin-K receptor, NKR, NK-3 receptor, NK-3R, Neurokinin B receptor, Tachykinin receptor 3, TACR3, NK3R, TAC3R

Target/Specificity

This TACR3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 406-435 amino acids from the C-terminal region of human TACR3.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TACR3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TACR3 Antibody (C-term) - Protein Information**Name** TACR3

Synonyms NK3R, TAC3R

Function This is a receptor for the tachykinin neuropeptide neuromedin-K (neurokinin B). It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system. The rank order of affinity of this receptor to tachykinins is: neuromedin-K > substance K > substance P.


Cellular Location

Cell membrane; Multi-pass membrane protein.

TACR3 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TACR3 Antibody (C-term) - Images

TACR3 Antibody (C-term) (Cat. #AP14222b) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the TACR3 antibody detected the TACR3 protein (arrow).

TACR3 Antibody (C-term) - Background

This gene belongs to a family of genes that function as receptors for tachykinins. Receptor affinities are specified by variations in the 5'-end of the sequence. The receptors belonging to this family are characterized by interactions with G proteins and 7 hydrophobic transmembrane regions. This gene encodes the receptor for the tachykinin neurokinin 3, also referred to as neurokinin B.

TACR3 Antibody (C-term) - References

Kalsi, G., et al. Hum. Mol. Genet. 19(12):2497-2506(2010)
Schuurhof, A., et al. Pediatr. Pulmonol. 45(6):608-613(2010)
Gianetti, E., et al. J. Clin. Endocrinol. Metab. 95(6):2857-2867(2010)
Young, J., et al. J. Clin. Endocrinol. Metab. 95(5):2287-2295(2010)
Fukami, M., et al. Horm Res Paediatr 73(6):477-481(2010)