

**Mouse Pdk3 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP14274b**

**Specification**

---

**Mouse Pdk3 Antibody (C-term) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">O922H2</a>
Other Accession	<a href="#">NP_663605.1</a>
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	387-414

**Mouse Pdk3 Antibody (C-term) - Additional Information**

**Gene ID** 236900

**Other Names**

[Pyruvate dehydrogenase (acetyl-transferring)] kinase isozyme 3, mitochondrial, Pyruvate dehydrogenase kinase isoform 3, Pdk3

**Target/Specificity**

This Mouse Pdk3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 387-414 amino acids from the C-terminal region of mouse Pdk3.

**Dilution**

WB~~1:1000

IHC-P~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Mouse Pdk3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**Mouse Pdk3 Antibody (C-term) - Protein Information**

**Name** Pdk3

**Function** Inhibits pyruvate dehydrogenase activity by phosphorylation of the E1 subunit PDHA1,

and thereby regulates glucose metabolism and aerobic respiration. Can also phosphorylate PDHA2. Decreases glucose utilization and increases fat metabolism in response to prolonged fasting, and as adaptation to a high-fat diet. Plays a role in glucose homeostasis and in maintaining normal blood glucose levels in function of nutrient levels and under starvation. Plays a role in the generation of reactive oxygen species (By similarity).


**Cellular Location**

Mitochondrion matrix.


**Mouse Pdk3 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Mouse Pdk3 Antibody (C-term) - Images**

Mouse Pdk3 Antibody (C-term) (Cat. #AP14274b) western blot analysis in mouse liver tissue lysates (35ug/lane). This demonstrates the Pdk3 antibody detected the Pdk3 protein (arrow).



Mouse Pdk3 Antibody (C-term) (AP14274b) immunohistochemistry analysis in formalin fixed and paraffin embedded mouse heart tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of Mouse Pdk3 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

#### **Mouse Pdk3 Antibody (C-term) - Background**

Pdk3 inhibits the mitochondrial pyruvate dehydrogenase complex by phosphorylation of the E1 alpha subunit, thus contributing to the regulation of glucose metabolism (By similarity).

#### **Mouse Pdk3 Antibody (C-term) - References**

Pagliarini, D.J., et al. Cell 134(1):112-123(2008)  
Blackshaw, S., et al. PLoS Biol. 2 (9), E247 (2004) :  
Visel, A., et al. Nucleic Acids Res. 32 (DATABASE ISSUE), D552-D556 (2004) :  
Mootha, V.K., et al. Cell 115(5):629-640(2003)

#### **Mouse Pdk3 Antibody (C-term) - Citations**

- [Metabolomic profiling analysis reveals chamber-dependent metabolite patterns in the mouse heart.](#)