

ING1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP14335a

Specification

ING1 Antibody (N-term) - Product Information

Application WB,E
Primary Accession Q9UK53

Other Accession NP 937862.1, NP 937861.1, NP 005528.3

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Human
Rabbit
Polyclonal
Rabbit IgG
A6738
1-30

ING1 Antibody (N-term) - Additional Information

Gene ID 3621

Other Names

Inhibitor of growth protein 1, ING1

Target/Specificity

This ING1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human ING1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ING1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ING1 Antibody (N-term) - Protein Information

Name ING1

Function Cooperates with p53/TP53 in the negative regulatory pathway of cell growth by modulating p53-dependent transcriptional activation. Implicated as a tumor suppressor gene.



Cellular Location Nucleus.

Tissue Location


Isoform 2 was expressed in all normal tissues and cells examined, as well as in all breast cancer and melanoma cell lines examined. Isoform 3 was expressed in testis, liver, and kidney, weakly expressed in colon and brain and not expressed in breast and cultured melanocytes. Isoform 4 was highly expressed in testis and weakly expressed in brain, but not expressed in breast, colon, kidney, melanocytes, breast cancer or melanoma cell lines

ING1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

ING1 Antibody (N-term) - Images



ING1 Antibody (N-term) (Cat. #AP14335a) western blot analysis in MDA-MB453 cell line lysates (35ug/lane). This demonstrates the ING1 antibody detected the ING1 protein (arrow).

ING1 Antibody (N-term) - Background

This gene encodes a tumor suppressor protein that can induce cell growth arrest and apoptosis. The encoded protein is a nuclear protein that physically interacts with the tumor suppressor protein TP53 and is a component of the p53 signaling pathway. Reduced expression and rearrangement of this gene have been detected in various cancers. Multiple alternatively spliced transcript variants encoding distinct isoforms have been reported.

ING1 Antibody (N-term) - References





Zhu, Z., et al. Hepatology 49(2):504-512(2009) Sayan, B., et al. Hybridoma (Larchmt) (2009) In press: Tallen, G., et al. Oncol. Res. 18 (2-3), 95-105 (2009): Halaschek-Wiener, J., et al. PLoS ONE 4 (8), E6641 (2009): Han, X., et al. Nat. Cell Biol. 10(11):1333-1340(2008)