

**OR4F16 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP14481a****Specification**

---

**OR4F16 Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q6IEY1</a>
Other Accession	<a href="#">O95013</a> , <a href="#">NP_001005221.2</a> , <a href="#">NP_001005224.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35074
Antigen Region	68-96

**OR4F16 Antibody (N-term) - Additional Information****Gene ID** 26683;729759;81399**Other Names**

Olfactory receptor 4F3/4F16/4F29, Olfactory receptor OR1-1, OR4F3

**Target/Specificity**

This OR4F16 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 68-96 amino acids from the N-terminal region of human OR4F16.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

OR4F16 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**OR4F16 Antibody (N-term) - Protein Information****Name** OR4F3**Function** Odorant receptor.


**Cellular Location**

Cell membrane; Multi-pass membrane protein.

**OR4F16 Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**OR4F16 Antibody (N-term) - Images**

OR4F16 Antibody (N-term) (Cat. #AP14481a) western blot analysis in Y79 cell line lysates (35ug/lane). This demonstrates the OR4F16 antibody detected the OR4F16 protein (arrow).

**OR4F16 Antibody (N-term) - Background**

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

**OR4F16 Antibody (N-term) - References**

Nusbaum, C., et al. Nature 439(7074):331-335(2006)

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)