

USP20 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14495C

Specification

USP20 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O9Y2K6
Other Accession	O8C6M1 , A7Z056 , NP_006667.3 , NP_001008563.2
Reactivity	Human, Mouse
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	102003
Antigen Region	310-339

USP20 Antibody (Center) - Additional Information

Gene ID 10868

Other Names

Ubiquitin carboxyl-terminal hydrolase 20, Deubiquitinating enzyme 20, Ubiquitin thioesterase 20, Ubiquitin-specific-processing protease 20, VHL-interacting deubiquitinating enzyme 2, hVDU2, USP20, KIAA1003, LSFR3A, VDU2

Target/Specificity

This USP20 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 310-339 amino acids from the Central region of human USP20.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

USP20 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

USP20 Antibody (Center) - Protein Information

Name USP20**Synonyms** KIAA1003, LSFR3A, VDU2

Function Deubiquitinating enzyme that plays a role in many cellular processes including autophagy, cellular antiviral response or membrane protein biogenesis (PubMed:[27801882](#), PubMed:[29487085](#)). Attenuates TLR4- mediated NF-kappa-B signaling by cooperating with beta-arrestin-2/ARRB2 and inhibiting TRAF6 autoubiquitination (PubMed:[26839314](#)). Promotes cellular antiviral responses by deconjugating 'Lys-33' and 'Lys-48'- linked ubiquitination of STING1 leading to its stabilization (PubMed:[27801882](#)). Plays an essential role in autophagy induction by regulating the ULK1 stability through deubiquitination of ULK1 (PubMed:[29487085](#)). Acts as a positive regulator for NF-kappa-B activation by TNF-alpha through deubiquitinating 'Lys-48'-linked polyubiquitination of SQSTM1, leading to its increased stability (PubMed:[32354117](#)). Acts as a regulator of G-protein coupled receptor (GPCR) signaling by mediating the deubiquitination beta-2 adrenergic receptor (ADRB2) (PubMed:[19424180](#)). Plays a central role in ADRB2 recycling and resensitization after prolonged agonist stimulation by constitutively binding ADRB2, mediating deubiquitination of ADRB2 and inhibiting lysosomal trafficking of ADRB2. Upon dissociation, it is probably transferred to the translocated beta-arrestins, possibly leading to beta-arrestins deubiquitination and disengagement from ADRB2 (PubMed:[19424180](#)). This suggests the existence of a dynamic exchange between the ADRB2 and beta-arrestins. Deubiquitinates DIO2, thereby regulating thyroid hormone regulation. Deubiquitinates HIF1A, leading to stabilize HIF1A and enhance HIF1A-mediated activity (PubMed:[15776016](#)). Deubiquitinates MCL1, a pivotal member of the anti- apoptotic Bcl-2 protein family to regulate its stability (PubMed:[35063767](#)). Within the endoplasmic reticulum, participates with USP33 in the rescue of post-translationally targeted membrane proteins that are inappropriately ubiquitinated by the cytosolic protein quality control in the cytosol (PubMed:[33792613](#)).


Cellular Location


Cytoplasm {ECO:0000250|UniProtKB:Q8C6M1}. Endoplasmic reticulum. Cytoplasm, perinuclear region. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome

USP20 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

USP20 Antibody (Center) - Images



USP20 Antibody (Center) (Cat. #AP14495c) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the USP20 antibody detected the USP20 protein (arrow).



USP20 Antibody (Center) (Cat. #AP14495c) western blot analysis in mouse testis tissue lysates (35ug/lane). This demonstrates the USP20 antibody detected the USP20 protein (arrow).

USP20 Antibody (Center) - Background

Deubiquitinating enzyme involved in beta-2 adrenergic receptor (ADRB2) recycling. Acts as a regulator of G-protein coupled receptor (GPCR) signaling by mediating the deubiquitination beta-2 adrenergic receptor (ADRB2). Plays a central role in ADRB2 recycling and resensitization after prolonged agonist stimulation by constitutively binding ADRB2, mediating deubiquitination of ADRB2 and inhibiting lysosomal trafficking of ADRB2. Upon dissociation, it is probably transferred to the translocated beta-arrestins, possibly leading to beta-arrestins deubiquitination and disengagement from ADRB2. This suggests the existence of a dynamic exchange between the ADRB2 and beta-arrestins. Deubiquitinates DIO2, thereby regulating thyroid hormone regulation. Deubiquitinates HIF1A, leading to stabilize HIF1A and enhance HIF1A-mediated activity. Mediates deubiquitination of both 'Lys-48'-and 'Lys-63'-linked polyubiquitin chains.

USP20 Antibody (Center) - References

Yoshida, T., et al. Int. J. Mol. Med. 25(4):649-656(2010)
Oguri, M., et al. Am. J. Hypertens. 23(1):70-77(2010)
Berthouze, M., et al. EMBO J. 28(12):1684-1696(2009)
Olsen, J.V., et al. Cell 127(3):635-648(2006)
Li, Z., et al. EMBO Rep. 6(4):373-378(2005)