

SEC23IP Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14511b

Specification

SEC23IP Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O9Y6Y8
Other Accession	NP_009121.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	724-752

SEC23IP Antibody (C-term) - Additional Information

Gene ID 11196

Other Names

SEC23-interacting protein, p125, SEC23IP

Target/Specificity

This SEC23IP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 724-752 amino acids from the C-terminal region of human SEC23IP.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SEC23IP Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SEC23IP Antibody (C-term) - Protein Information

Name SEC23IP

Function Plays a role in the organization of endoplasmic reticulum exit sites. Specifically binds to phosphatidylinositol 3-phosphate (PI(3)P), phosphatidylinositol 4-phosphate (PI(4)P) and phosphatidylinositol 5-phosphate (PI(5)P).

Cellular Location

Cytoplasmic vesicle, COPII-coated vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Endoplasmic reticulum


Tissue Location

Ubiquitously expressed with stronger levels detected in heart, liver and skeletal muscle

SEC23IP Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SEC23IP Antibody (C-term) - Images

SEC23IP Antibody (C-term) (Cat. #AP14511b) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the SEC23IP antibody detected the SEC23IP protein (arrow).

SEC23IP Antibody (C-term) - Background

COPII-coated vesicles are involved in protein transport from the endoplasmic reticulum to the Golgi apparatus. The protein encoded by this gene was identified by its interaction with a mouse protein similar to yeast Sec23p, an essential component of the COPII. This protein shares significant similarity with phospholipid-modifying proteins, especially phosphatidic acid preferring-phospholipase A1. Overexpression of this protein has been shown to cause disorganization of the endoplasmic reticulum-Golgi intermediate compartment and Golgi apparatus, which suggests its role in the early secretory pathway. [provided by RefSeq].

SEC23IP Antibody (C-term) - References

Ong, Y.S., et al. J. Cell Biol. 190(3):331-345(2010)
Trynka, G., et al. Gut 58(8):1078-1083(2009)
Li, H., et al. J. Biol. Chem. 281(21):14748-14755(2006)
Grupe, A., et al. Am. J. Hum. Genet. 78(1):78-88(2006)
Shimoi, W., et al. J. Biol. Chem. 280(11):10141-10148(2005)

SEC23IP Antibody (C-term) - Citations

- [A cascade of ER exit site assembly that is regulated by p125A and lipid signals.](#)