

**IRX3 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP14549a**

**Specification**

---

**IRX3 Antibody (N-term) - Product Information**

Application	IHC-P, WB,E
Primary Accession	<a href="#">P78415</a>
Other Accession	<a href="#">O42261</a> , <a href="#">P81067</a> , <a href="#">NP_077312.2</a>
Reactivity	Human
Predicted	Mouse, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	52119
Antigen Region	111-140

**IRX3 Antibody (N-term) - Additional Information**

**Gene ID** 79191

**Other Names**

Iroquois-class homeodomain protein IRX-3, Homeodomain protein IRXB1, Iroquois homeobox protein 3, IRX3, IRXB1

**Target/Specificity**

This IRX3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 111-140 amino acids from the N-terminal region of human IRX3.

**Dilution**

IHC-P~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

IRX3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**IRX3 Antibody (N-term) - Protein Information**

**Name** IRX3

**Synonyms** IRXB1

**Function** Transcription factor involved in SHH-dependent neural patterning. Together with NKX2-2 and NKX6-1 acts to restrict the generation of motor neurons to the appropriate region of the neural tube. Belongs to the class I proteins of neuronal progenitor factors, which are repressed by SHH signals. Involved in the transcriptional repression of MNX1 in non-motor neuron cells. Acts as a regulator of energy metabolism.

**Cellular Location**


Nucleus.

### IRX3 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### IRX3 Antibody (N-term) - Images



Anti-IRX3 Antibody (N-term) at 1:1000 dilution + SH-SY5Y whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 52 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



IRX3 Antibody (N-term) (AP14549a) immunohistochemistry analysis in formalin fixed and paraffin embedded human breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of IRX3 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

#### **IRX3 Antibody (N-term) - Background**

IRX3 is a member of the Iroquois homeobox gene family (see IRX1; MIM 606197) and plays a role in an early step of neural development (Bellefroid et al., 1998 [PubMed 9427753]). Members of this family appear to play multiple roles during pattern formation of vertebrate embryos (Lewis et al., 1999 [PubMed 10370142]).

#### **IRX3 Antibody (N-term) - References**

Ragvin, A., et al. Proc. Natl. Acad. Sci. U.S.A. 107(2):775-780(2010)  
Trynka, G., et al. Gut 58(8):1078-1083(2009)  
Lewis, M.T., et al. Cell Tissue Res. 296(3):549-554(1999)  
Bellefroid, E.J., et al. EMBO J. 17(1):191-203(1998)