

Mouse Epha2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14619c**Specification**

Mouse Epha2 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q03145
Other Accession	NP_034269.2
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	108852
Antigen Region	296-323

Mouse Epha2 Antibody (Center) - Additional Information**Gene ID** 13836**Other Names**

Ephrin type-A receptor 2, Epithelial cell kinase, Tyrosine-protein kinase receptor ECK, Tyrosine-protein kinase receptor MPK-5, Tyrosine-protein kinase receptor SEK-2, Epha2, Eck, Myk2, Sek2

Target/Specificity

This Mouse Epha2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 296-323 amino acids from the Central region of mouse Epha2.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Epha2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Epha2 Antibody (Center) - Protein Information**Name** Epha2

Synonyms Eck, Myk2, Sek2

Function Receptor tyrosine kinase which binds promiscuously membrane- bound ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Activated by the ligand ephrin- A1/EFNA1 regulates migration, integrin-mediated adhesion, proliferation and differentiation of cells (PubMed:[29749928](#)). Regulates cell adhesion and differentiation through DSG1/desmoglein-1 and inhibition of the ERK1/ERK2 signaling pathway. May also participate in UV radiation- induced apoptosis and have a ligand-independent stimulatory effect on chemotactic cell migration. During development, may function in distinctive aspects of pattern formation and subsequently in development of several fetal tissues. Involved for instance in angiogenesis, in early hindbrain development and epithelial proliferation and branching morphogenesis during mammary gland development. Engaged by the ligand ephrin-A5/EFNA5 may regulate lens fiber cells shape and interactions and be important for lens transparency development and maintenance. With ephrin-A2/EFNA2 may play a role in bone remodeling through regulation of osteoclastogenesis and osteoblastogenesis.

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:P29317}; Single-pass type I membrane protein. Cell projection, ruffle membrane {ECO:0000250|UniProtKB:P29317}; Single-pass type I membrane protein. Cell projection, lamellipodium membrane {ECO:0000250|UniProtKB:P29317}; Single-pass type I membrane protein. Cell junction, focal adhesion {ECO:0000250|UniProtKB:P29317}. Note=Present at regions of cell-cell contacts but also at the leading edge of migrating cells. Relocates from the plasma membrane to the cytoplasmic and perinuclear regions in cancer cells. {ECO:0000250|UniProtKB:P29317}


Tissue Location


Expressed in the lung, intestine and liver (PubMed:11287184). Expressed in myogenic progenitor cells (PubMed:27446912).

Mouse EphA2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mouse EphA2 Antibody (Center) - Images



Anti-Mouse Epha2 Antibody (Center) at 1:1000 dilution + mouse lung lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 109 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Mouse Epha2 Antibody (Center) - Background

Receptor for members of the ephrin-A family. Binds to ephrin-A1, -A3, -A4 and -A5. Induces apoptosis in a TP53/p53-independent, caspase-8-dependent manner (By similarity). Plays an important role in angiogenesis and tumor neovascularization. The recruitment of VAV2, VAV3 and PI3-kinase p85 subunit by phosphorylated EPHA2 is critical for EFNA1-induced RAC1 GTPase activation and vascular endothelial cell migration and assembly. May function in distinctive aspects of pattern formation and subsequently in development of several fetal tissues. May be involved in cell-cell interactions guiding early hindbrain development.

Mouse Epha2 Antibody (Center) - References

Islam, S., et al. Dig. Dis. Sci. 55(9):2478-2488(2010)
Kim, J., et al. Mol. Cell. Biol. 30(7):1582-1592(2010)
Zirzow, S., et al. Dev. Biol. 336(2):145-155(2009)
Jun, G., et al. PLoS Genet. 5 (7), E1000584 (2009) :
North, H.A., et al. Development 136(14):2467-2476(2009)