

SNX6 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1464a

Specification

SNX6 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q9UNH7
Other Accession	Q6P8X1
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	46649
Antigen Region	81-110

SNX6 Antibody (N-term) - Additional Information

Gene ID 58533

Other Names

Sorting nexin-6, TRAF4-associated factor 2, Sorting nexin-6, N-terminally processed, SNX6

Target/Specificity

This SNX6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 81-110 amino acids from the N-terminal region of human SNX6.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SNX6 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SNX6 Antibody (N-term) - Protein Information

Name SNX6

Function Involved in several stages of intracellular trafficking. Interacts with membranes phosphatidylinositol 3,4-bisphosphate and/or phosphatidylinositol 4,5-bisphosphate (Probable). Acts in part as component of the retromer membrane-deforming SNX-BAR subcomplex (PubMed:[19935774](#)). The SNX-BAR retromer mediates retrograde transport of cargo proteins from endosomes to the trans-Golgi network (TGN) and is involved in endosome-to-plasma membrane transport for cargo protein recycling. The SNX-BAR subcomplex functions to deform the donor membrane into a tubular profile called endosome-to-TGN transport carrier (ETC) (Probable). Does not have in vitro vesicle-to-membrane remodeling activity (PubMed:[23085988](#)). Involved in retrograde endosome- to-TGN transport of lysosomal enzyme receptor IGF2R (PubMed:[17148574](#)). May function as link between transport vesicles and dynactin (Probable). Negatively regulates retrograde transport of BACE1 from the cell surface to the trans-Golgi network (PubMed:[20354142](#)). Involved in E-cadherin sorting and degradation; inhibits PIP5K1C isoform 3-mediated E-cadherin degradation (PubMed:[24610942](#)). In association with GIT1 involved in EGFR degradation. Promotes lysosomal degradation of CDKN1B (By similarity). May contribute to transcription regulation (Probable).

Cellular Location


Early endosome. Early endosome membrane; Peripheral membrane protein; Cytoplasmic side Cytoplasmic vesicle. Cytoplasm. Nucleus. Note=Interaction with SNX1 or SNX2 promotes location at endosome membranes (PubMed:[19935774](#)). Only a minor proportion is seen in the nucleus.

SNX6 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SNX6 Antibody (N-term) - Images



Western blot analysis of SNX6 Antibody (N-term) (Cat# AP1464a) in HepG2 cell line lysates (35ug/lane). SNX6 (arrow) was detected using the purified Pab.

SNX6 Antibody (N-term) - Background

SNX6 interacts with members of the transforming growth factor-beta family of receptor serine-threonine kinases. These receptors belong to two classes: type II receptors that bind ligand, and type I receptors that are subsequently recruited to transduce the signal. Of the type II receptors, SNX6 was found to interact strongly with ActRIIB and more moderately with wild type and kinase-defective mutants of TbetaRII. Of the type I receptors, SNX6 was found to interact only with inactivated TbetaRI. SNXs 1-4 also interacted with the transforming growth factor-beta receptor family, showing different receptor preferences. Conversely, SNX6 behaved similarly to the other SNX proteins in its interactions with receptor tyrosine kinases. Strong heteromeric interactions were also seen among SNX1, -2, -4, and -6, suggesting the formation *in vivo* of oligomeric complexes. These findings are the first evidence for the association of the SNX family of molecules with receptor serine-threonine kinases.

SNX6 Antibody (N-term) - References

Parks W.T., J. Biol. Chem. 276:19332-19339(2001).