

**CD274 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP14682b**

**Specification**

---

**CD274 Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O9NZQ7</a>
Other Accession	<a href="#">NP_054862.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	261-290

**CD274 Antibody (C-term) - Additional Information**

**Gene ID** 29126

**Other Names**

Programmed cell death 1 ligand 1, PD-L1, PDCD1 ligand 1, Programmed death ligand 1, B7 homolog 1, B7-H1, CD274, CD274, B7H1, PDCD1L1, PDCD1LG1, PDL1

**Target/Specificity**

This CD274 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 261-290 amino acids from the C-terminal region of human CD274.

**Dilution**

WB~~1:2000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CD274 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CD274 Antibody (C-term) - Protein Information**

**Name** CD274 ([HGNC:17635](#))

**Function** Plays a critical role in induction and maintenance of immune tolerance to self

(PubMed:[11015443](#), PubMed:[28813410](#), PubMed:[28813417](#), PubMed:[31399419](#)). As a ligand for the inhibitory receptor PDCD1/PD-1, modulates the activation threshold of T-cells and limits T-cell effector response (PubMed:[11015443](#), PubMed:[28813410](#), PubMed:[28813417](#), PubMed:[36727298](#)). Through a yet unknown activating receptor, may costimulate T-cell subsets that predominantly produce interleukin-10 (IL10) (PubMed:[10581077](#)). Can also act as a transcription coactivator: in response to hypoxia, translocates into the nucleus via its interaction with phosphorylated STAT3 and promotes transcription of GSDMC, leading to pyroptosis (PubMed:[32929201](#)).

#### Cellular Location

Cell membrane; Single-pass type I membrane protein. Early endosome membrane; Single-pass type I membrane protein. Recycling endosome membrane; Single-pass type I membrane protein. Nucleus. Note=Associates with CMTM6 at recycling endosomes, where it is protected from being targeted for lysosomal degradation (PubMed:28813417). Translocates to the nucleus in response to hypoxia via its interaction with phosphorylated STAT3 (PubMed:32929201). [Isoform 2]: Endomembrane system; Single-pass type I membrane protein

#### Tissue Location


Highly expressed in the heart, skeletal muscle, placenta and lung. Weakly expressed in the thymus, spleen, kidney and liver. Expressed on activated T- and B-cells, dendritic cells, keratinocytes and monocytes.

#### CD274 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### CD274 Antibody (C-term) - Images



Anti-CD274 Antibody (C-term) at 1:2000 dilution + A549 whole cell lysate Lysates/proteins at 20

µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 33 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



All lanes : Anti-CD274 Antibody (C-term) at 1:2000 dilution Lane 1: A549 whole cell lysate Lane 2: HCT116 whole cell lysate Lane 3: HeLa whole cell lysate Lane 4: Raji whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 33 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

#### CD274 Antibody (C-term) - Background

Involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PDCD1-independent manner. Interaction with PDCD1 inhibits T-cell proliferation and cytokine production.

#### CD274 Antibody (C-term) - References

- Berthon, C., et al. Cancer Immunol. Immunother. 59(12):1839-1849(2010)
- Dianzani, C., et al. J. Immunol. 185(7):3970-3979(2010)
- Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)
- Alvarez, I.B., et al. J. Infect. Dis. 202(4):524-532(2010)
- Francisco, L.M., et al. Immunol. Rev. 236, 219-242 (2010) :