

ZNF398 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14859c

Specification

ZNF398 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q8TD17
Other Accession	NP_065832.1 , NP_733787.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	71311
Antigen Region	186-215

ZNF398 Antibody (Center) - Additional Information

Gene ID 57541

Other Names

Zinc finger protein 398, Zinc finger DNA-binding protein p52/p71, ZNF398, KIAA1339, ZER6

Target/Specificity

This ZNF398 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 186-215 amino acids from the Central region of human ZNF398.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ZNF398 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF398 Antibody (Center) - Protein Information

Name ZNF398

Synonyms KIAA1339, ZER6

Function Functions as a transcriptional activator.

Cellular Location


Nucleus.

ZNF398 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF398 Antibody (Center) - Images



ZNF398 Antibody (Center) (Cat. #AP14859c) western blot analysis in SK-BR-3 cell line lysates (35ug/lane). This demonstrates the ZNF398 antibody detected the ZNF398 protein (arrow).

ZNF398 Antibody (Center) - Background

This gene encodes a member of the Kruppel family of C2H2-type zinc-finger transcription factor proteins. The encoded protein acts as a transcriptional activator. Two transcript variants encoding distinct isoforms have been identified for this gene. Other transcript variants have been described, but their full length sequence has not been determined.

ZNF398 Antibody (Center) - References

Conroy, A.T., et al. J. Biol. Chem. 277(11):9326-9334(2002)