

Mouse Camkk2 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP14951a

Specification

Mouse Camkk2 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	<u>Q8C078</u>
Other Accession	<u>NP_663333.1</u>
Reactivity	Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	43-71

Mouse Camkk2 Antibody (N-term) - Additional Information

Gene ID 207565

Other Names

Calcium/calmodulin-dependent protein kinase kinase 2, CaM-KK 2, CaM-kinase kinase 2, CaMKK 2, Calcium/calmodulin-dependent protein kinase kinase beta, CaM-KK beta, CaM-kinase kinase beta, CaMKK beta, Camkk2, Kiaa0787

Target/Specificity

This Mouse Camkk2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 43-71 amino acids from the N-terminal region of mouse Camkk2.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Camkk2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Camkk2 Antibody (N-term) - Protein Information

Name Camkk2



Synonyms Kiaa0787

Function Calcium/calmodulin-dependent protein kinase belonging to a proposed calcium-triggered signaling cascade involved in a number of cellular processes. Phosphorylates CAMK1, CAMK4 and CAMK1D (By similarity). Efficiently phosphorylates 5'-AMP-activated protein kinase (AMPK) trimer, including that consisting of PRKAA1, PRKAB1 and PRKAG1. This phosphorylation is stimulated in response to Ca(2+) signals (By similarity). May play a role in neurite growth. Isoform 2 may promote neurite elongation, while isoform 1 may promoter neurite branching (By similarity). May be involved in hippocampal activation of CREB1.

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q96RR4}. Cytoplasm {ECO:0000250|UniProtKB:Q96RR4}. Cell projection, neuron projection {ECO:0000250|UniProtKB:Q96RR4}. Note=Predominantly nuclear in unstimulated cells, relocalizes into cytoplasm and neurites after forskolin induction. {ECO:0000250|UniProtKB:Q96RR4}

Tissue Location

Expressed in all tissues tested. A differential expression pattern compared to CAMKK1 is observed in the brain

Mouse Camkk2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- <u>Dot Blot</u>
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Mouse Camkk2 Antibody (N-term) - Images



Anti-Mouse Camkk2 Antibody (N-term) at 1:2000 dilution + mouse liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000



dilution. Predicted band size : 65 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Mouse Camkk2 Antibody (N-term) - Background

Calcium/calmodulin-dependent protein kinase belonging to a proposed calcium-triggered signaling cascade involved in a number of cellular processes. Phosphorylates CAMK1, CAMK4 and CAMK1D (By similarity). Seems to be involved in hippocampal activation of CREB1.

Mouse Camkk2 Antibody (N-term) - References

Jin, X.L., et al. Biol. Reprod. 82(2):459-468(2010) Kokubo, M., et al. J. Neurosci. 29(28):8901-8913(2009) Anderson, K.A., et al. Cell Metab. 7(5):377-388(2008) Park, C.S., et al. Neuroscience 151(1):43-55(2008) Hoyer-Hansen, M., et al. Mol. Cell 25(2):193-205(2007)