

**DKK4 Antibody (N-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP1524a****Specification**

---

**DKK4 Antibody (N-term) - Product Information**

Application	IHC-P, WB,E
Primary Accession	<a href="#">Q9UBT3</a>
Other Accession	<a href="#">NP_055235</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	1-30

**DKK4 Antibody (N-term) - Additional Information****Gene ID** 27121**Other Names**

Dickkopf-related protein 4, Dickkopf-4, Dkk-4, hDkk-4, Dickkopf-related protein 4 short form, DKK4

**Target/Specificity**

This DKK4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human DKK4.

**Dilution**

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

DKK4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**DKK4 Antibody (N-term) - Protein Information****Name** DKK4**Function** Antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt and by

forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6. DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as antero-posterior axial patterning, limb development, somitogenesis and eye formation. In the adult, Dkks are implicated in bone formation and bone disease, cancer and Alzheimer disease (By similarity).

**Cellular Location**

Secreted.


**Tissue Location**

Expressed in cerebellum, T-cells, esophagus and lung


**DKK4 Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**DKK4 Antibody (N-term) - Images**

The anti-DKK4 (N-term) (Cat. #AP1524a) is used in Western blot to detect DKK4 in A375 cell lysate.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with DKK4 antibody (N-term) (Cat.#AP1524a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

#### **DKK4 Antibody (N-term) - Background**

DKK4, like DKK1, DKK2, and DKK3, possesses an N-terminal signal peptide and 2 conserved cysteine-rich domains, which are separated by a linker region and contain 10 cysteine residues each. The second cysteine region has a putative lipid-binding function that may facilitate WNT/DKK interactions at the plasma membrane. The linker region contains 50 to 55 amino acids in DKK1, DKK2, and DKK4, whereas in DKK3 it contains only 12 amino acids. All DKKs have several potential sites for cleavage by furin-type proteases. Northern blot analysis detected no expression of DKK4, but RT-PCR analysis detected DKK4 expression in cerebellum, T-cell, esophagus, and lung cDNA libraries. DKK4 blocks *Xenopus* Wnt8, Wnt3a, and Wnt2b, but not Dsh or Fz8, induction of a secondary axis in frog embryos, indicating that DKKs antagonize WNT function upstream of WNT receptors.

#### **DKK4 Antibody (N-term) - References**

Krupnik, V.E., et al., *Gene* 238(2):301-313 (1999). Yoshida, S., et al., *Cytogenet. Cell Genet.* 94 (1-2), 88-89 (2001).

#### **DKK4 Antibody (N-term) - Citations**

- [SRC-2-mediated coactivation of anti-tumorigenic target genes suppresses MYC-induced liver cancer.](#)
- [Dickkopf 4 \(DKK4\) acts on Wnt/ \$\beta^2\$ -catenin pathway by influencing  \$\beta^2\$ -catenin in hepatocellular carcinoma.](#)
- [DICKKOPF-4 activates the noncanonical c-Jun-NH2 kinase signaling pathway while inhibiting the Wnt-canonical pathway in human renal cell carcinoma.](#)
- [DICKKOPF-4 and -2 genes are upregulated in human colorectal cancer.](#)