

Connexin 50 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1548a**Specification**

Connexin 50 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P48165
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	99-130

Connexin 50 Antibody (N-term) - Additional Information**Gene ID** 2703**Other Names**

Gap junction alpha-8 protein, Connexin-50, Cx50, Lens fiber protein MP70, GJA8

Target/Specificity

This Connexin 50 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 99-130 amino acids from the N-terminal region of human Connexin 50.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Connexin 50 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Connexin 50 Antibody (N-term) - Protein Information**Name** GJA8

Function Structural component of eye lens gap junctions (PubMed:[18006672](#), PubMed:[19756179](#)). Gap junctions are dodecameric channels that connect the cytoplasm of adjoining cells. They are formed by the docking of two hexameric hemichannels, one from each cell membrane (By similarity). Small molecules and ions diffuse from one cell to a neighboring cell via the central pore

(PubMed:[18006672](#), PubMed:[19756179](#)).

Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P55917}. Cell junction, gap junction

Tissue Location


Eye lens..

Connexin 50 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Connexin 50 Antibody (N-term) - Images



Western blot analysis of anti-GJA8 Antibody (N-term)(Cat.#AP1548a) in HepG2 cell line lysates (35ug/lane). GJA8(arrow) was detected using the purified Pab.

Connexin 50 Antibody (N-term) - Background

GJA8 is a an integral membrane protein that belongs to the connexin family, alpha-type (group II) subfamily. One gap junction consists of a cluster of closely packed pairs of transmembrane channels, the connexons, through which materials of low MW diffuse from one cell to a neighboring cell. A connexon is composed of a hexamer of connexins. This particular connexin only forms junctional channels. GJA8 is expressed in the eye lens, and defects in GJA8 are the cause of zonular pulverulent cataract type 1 (CZP1), a form of autosomal dominant congenital cataract.

Connexin 50 Antibody (N-term) - References

Shiels, A., et al., Am. J. Hum. Genet. 62(3):526-532 (1998).

Church, R.L., et al., Curr. Eye Res. 14(10):979-981 (1995).

Church, R.L., et al., Curr. Eye Res. 14(3):215-221 (1995).

Connexin 50 Antibody (N-term) - Citations

- [Intramolecular loop/tail interactions are essential for connexin 43-hemichannel activity.](#)