

KANK2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16024c

Specification

KANK2 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q63ZY3
Other Accession	NP_056308.3 , NP_001129663.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	91174
Antigen Region	465-493

KANK2 Antibody (Center) - Additional Information

Gene ID 25959

Other Names

KN motif and ankyrin repeat domain-containing protein 2, Ankyrin repeat domain-containing protein 25, Matrix-remodeling-associated protein 3, SRC-1-interacting protein, SIP, SRC-interacting protein, SRC1-interacting protein, KANK2, ANKRD25, KIAA1518, MXRA3, SIP

Target/Specificity

This KANK2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 465-493 amino acids from the Central region of human KANK2.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

KANK2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

KANK2 Antibody (Center) - Protein Information

Name KANK2

Synonyms ANKRD25, KIAA1518, MXRA3, SIP

Function Involved in transcription regulation by sequestering in the cytoplasm nuclear receptor coactivators such as NCOA1, NCOA2 and NCOA3 (PubMed:[17476305](#)). Involved in regulation of caspase-independent apoptosis by sequestering the proapoptotic factor AIFM1 in mitochondria (PubMed:[22371500](#)). Pro-apoptotic stimuli can induce its proteasomal degradation allowing the translocation of AIFM1 to the nucleus to induce apoptosis (PubMed:[22371500](#)). Involved in the negative control of vitamin D receptor signaling pathway (PubMed:[24671081](#)). Involved in actin stress fibers formation through its interaction with ARHGDI and the regulation of the Rho signaling pathway (PubMed:[17996375](#), PubMed:[25961457](#)). May thereby play a role in cell adhesion and migration, regulating for instance podocytes migration during development of the kidney (PubMed:[25961457](#)). Through the Rho signaling pathway may also regulate cell proliferation (By similarity).

Cellular Location

Cytoplasm. Mitochondrion


Tissue Location

Strongly expressed in cervix, colon, heart, kidney and lung. Expressed in kidney glomerular podocytes and mesangial cells (at protein level).

KANK2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KANK2 Antibody (Center) - Images

KANK2 Antibody (Center) (Cat. #AP16024c) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the KANK2 antibody detected the KANK2 protein (arrow).

KANK2 Antibody (Center) - Background

ANKRD25 contains 5 ANK repeats. It is strongly expressed in cervix, colon, heart, kidney and lung.

KANK2 Antibody (Center) - References

Zhu, Y., et al. *Biochim. Biophys. Acta* 1780(2):128-133(2008)

Zhang, Y., et al. *EMBO J.* 26(11):2645-2657(2007)

Olsen, J.V., et al. *Cell* 127(3):635-648(2006)

Wistow, G., et al. *Mol. Vis.* 8, 205-220 (2002) :