

Mouse Cdk18 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP16070b

Specification

Mouse Cdk18 Antibody (C-term) - Product Information

Application WB,E
Primary Accession O04899

Other Accession <u>035832</u>, <u>NP_032821.1</u>

Reactivity
Predicted
Rat
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Mouse
Rat
Rabbit
Rabbit
Polyclonal
Rabbit IgG
51848
422-449

Mouse Cdk18 Antibody (C-term) - Additional Information

Gene ID 18557

Other Names

Cyclin-dependent kinase 18, Cell division protein kinase 18, PCTAIRE-motif protein kinase 3, Serine/threonine-protein kinase PCTAIRE-3, Cdk18, Pctaire3, Pctk3

Target/Specificity

This Mouse Cdk18 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 422-449 amino acids from the C-terminal region of mouse Cdk18.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Cdk18 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Cdk18 Antibody (C-term) - Protein Information

Name Cdk18



Synonyms Pctaire3, Pctk3

Function May play a role in signal transduction cascades in terminally differentiated cells.

Tissue Location


In brain, kidney, intestine and at a much lower level, in fetal tissues

Mouse Cdk18 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Mouse Cdk18 Antibody (C-term) - Images



Mouse Cdk18 Antibody (C-term) (Cat. #AP16070b) western blot analysis in mouse stomach tissue lysates (35ug/lane). This demonstrates the Cdk18 antibody detected the Cdk18 protein (arrow).

Mouse Cdk18 Antibody (C-term) - Background

Cdk18 may play a role in signal transduction cascades in terminally differentiated cells.

Mouse Cdk18 Antibody (C-term) - References

Zambrowicz, B.P., et al. Proc. Natl. Acad. Sci. U.S.A. 100(24):14109-14114(2003) Rhee, K., et al. Dev. Dyn. 204(4):406-420(1995) Okuda, T., et al. Genomics 21(1):217-221(1994) Okuda, T., et al. Oncogene 7(11):2249-2258(1992)