

RP13-102H20.1/ARHGAP36 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP16204b

Specification

RP13-102H20.1/ARHGAP36 Antibody (C-term) - Product Information

WB,E Application **Primary Accession 06ZRI8** Other Accession NP 659404.2 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 61664 Antigen Region 13-102

RP13-102H20.1/ARHGAP36 Antibody (C-term) - Additional Information

Gene ID 158763

Other Names

Rho GTPase-activating protein 36, ARHGAP36

Target/Specificity

This RP13-102H20.1/ARHGAP36 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 373-402 amino acids from the C-terminal region of human RP13-102H20.1/ARHGAP36.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RP13-102H20.1/ARHGAP36 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RP13-102H20.1/ARHGAP36 Antibody (C-term) - Protein Information

Name ARHGAP36

Function GTPase activator for the Rho-type GTPases by converting them to an inactive



GDP-bound state.

Tissue Location


Detected in the outer root sheath of hair follicles at the level of the stem cell bulge, during the anagen and telogen phases of hair growth (at protein level)

RP13-102H20.1/ARHGAP36 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

RP13-102H20.1/ARHGAP36 Antibody (C-term) - Images



RP13-102H20.1/ARHGAP36 Antibody (C-term) (Cat. #AP16204b) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the RP13-102H20.1 antibody detected the RP13-102H20.1 protein (arrow).

RP13-102H20.1/ARHGAP36 Antibody (C-term) - Background

GTPase activator for the Rho-type GTPases by converting them to an inactive GDP-bound state (By similarity).

RP13-102H20.1/ARHGAP36 Antibody (C-term) - References

Ross, M.T., et al. Nature 434(7031):325-337(2005)